

SEMIDISCRETE FINITE ELEMENT APPROXIMATIONS OF A LINEAR FLUID-STRUCTURE INTERACTION PROBLEM*

Q. DU[†], M. D. GUNZBURGER[‡], L. S. HOU[§], AND J. LEE[¶]

Abstract. Semidiscrete finite element approximations of a linear fluid-structure interaction problem are studied. First, results concerning a divergence-free weak formulation of the interaction problem are reviewed. Next, semidiscrete finite element approximations are defined, and the existence of finite element solutions is proved with the help of an auxiliary, discretely divergence-free formulation. A discrete inf-sup condition is verified, and the existence of a finite element pressure is established. Strong a priori estimates for the finite element solutions are also derived. Then, by passing to the limit in the finite element approximations, the existence of a strong solution is demonstrated and semidiscrete error estimates are obtained.

Key words. fluid-structure interactions, finite element methods, error estimates

AMS subject classifications. 65M60, 76M10, 76D07, 73V05, 73C02

DOI. 10.1137/S0036142903408654

1. Introduction. Fluid-structure interaction problems have been extensively studied in recent years both analytically and computationally. The book [28] and the special issue [30] give accounts of the state of the art from the engineering points of view. In addition, a short discussion of the literature can be found in [10]. The references in [10] include [4, 18, 29] for fluid-structure interactions involving elementary fluids, [2, 3, 32] for fluid-structure interactions involving inviscid fluids, and [6, 7, 8, 9, 11, 13, 14, 15, 16, 17, 20, 21, 22, 26, 27, 33, 34] for interactions between viscous, incompressible fluids and elastic solids.

In [10], we analyzed a model for the interactions between Stokesian fluids and linear elastic solids. This paper is devoted to the finite element analysis of that model. As in [10], we assume that the fluid and solid occupy two adjacent open Lipschitz domains, $\Omega_1 \subset \mathbb{R}^d$ and $\Omega_2 \subset \mathbb{R}^d$, respectively, where $d = 2$ or 3 is the space dimension. We denote by Ω the entire fluid-solid region under consideration; i.e., Ω is the interior of $\overline{\Omega}_1 \cup \overline{\Omega}_2$. Let $\Gamma_0 = \partial\Omega_1 \cap \partial\Omega_2$ denote the interface between the fluid and solid, and let $\Gamma_1 = \partial\Omega_1 \setminus \Gamma_0$ and $\Gamma_2 = \partial\Omega_2 \setminus \Gamma_0$ denote the parts of the fluid and solid boundaries, respectively, excluding the interface Γ_0 . For obvious reasons we assume that $\text{meas}(\Gamma_1 \cup \Gamma_2) \neq 0$.

*Received by the editors May 31, 2002; accepted for publication (in revised form) March 7, 2003; published electronically January 6, 2004.

<http://www.siam.org/journals/sinum/42-1/40865.html>

[†]Department of Mathematics, Penn State University, State College, PA 16802 (qdu@math.psu.edu). The work of this author was supported in part by the National Science Foundation under grant DMS-0196522.

[‡]School of Computational Science and Information Technology, Florida State University, Tallahassee, FL 32306-4120 (gunzburg@csit.fsu.edu). The work of this author was supported in part by the National Science Foundation under grant DMS-9806358.

[§]Department of Mathematics, Iowa State University, Ames, IA 50011-2064 (hou@math.iastate.edu).

[¶]Department of Mathematics, Carnegie Mellon University, Pittsburgh, PA 15213 (jeehyun@andrew.cmu.edu).

In the fluid region Ω_1 , we apply the Stokes system

$$(1.1) \quad \left\{ \begin{array}{ll} \rho_1 \mathbf{v}_t + \nabla p - \mu_1 \nabla \cdot (\nabla \mathbf{v} + \nabla \mathbf{v}^T) = \rho_1 \mathbf{f}_1 & \text{in } \Omega_1, \\ \nabla \cdot \mathbf{v} = 0 & \text{in } \Omega_1, \\ \mathbf{v} = 0 & \text{on } \Gamma_1, \\ \mathbf{v}|_{t=0} = \mathbf{v}_0 & \text{in } \Omega_1, \end{array} \right.$$

where \mathbf{v} denotes the fluid velocity, p the fluid pressure, \mathbf{f}_1 the given body force per unit mass, ρ_1 and μ_1 the constant fluid density and viscosity, and \mathbf{v}_0 the given initial velocity.

In the solid region, we apply the equations of linear elasticity

$$(1.2) \quad \left\{ \begin{array}{ll} \rho_2 \mathbf{u}_{tt} - \mu_2 \nabla \cdot (\nabla \mathbf{u} + \nabla \mathbf{u}^T) - \lambda_2 \nabla (\nabla \cdot \mathbf{u}) = \rho_2 \mathbf{f}_2 & \text{in } \Omega_2, \\ \mathbf{u} = 0 & \text{on } \Gamma_2, \\ \mathbf{u}|_{t=0} = \mathbf{u}_0 \quad \text{and} \quad \mathbf{u}_t|_{t=0} = \mathbf{u}_1 & \text{in } \Omega_2, \end{array} \right.$$

where \mathbf{u} denotes the displacement of the solid, \mathbf{f}_2 the given loading force per unit mass, μ_2 and λ_2 the Lamé constants, ρ_2 the constant solid density, and \mathbf{u}_0 and \mathbf{u}_1 the given initial data.

Across the *fixed* interface Γ_0 between the fluid and solid, the velocity and stress vector are continuous. Thus, we have

$$(1.3) \quad \mathbf{u}_t = \mathbf{v} \quad \text{on } \Gamma_0$$

and

$$(1.4) \quad \mu_2 (\nabla \mathbf{u} + \nabla \mathbf{u}^T) \cdot \mathbf{n}_2 + \lambda_2 (\nabla \cdot \mathbf{u}) \mathbf{n}_2 = p \mathbf{n}_1 - \mu_1 (\nabla \mathbf{v} + \nabla \mathbf{v}^T) \cdot \mathbf{n}_1 \quad \text{on } \Gamma_0,$$

where \mathbf{n}_i is the outward-pointing unit normal vector along $\partial\Omega_i$, $i = 1, 2$.

The physical validity of the model (1.1)–(1.4) was explained in [10]. Previous work concerning this model include, as cited in [10], eigenmode analysis [34], homogenization [8], the one-dimensional case [11], and a numerical algorithm [13]. In [10], weak formulations for (1.1)–(1.4) were defined, and the existence of weak solutions was established. The proof for the existence result was based on Galerkin approximations using divergence-free basis functions, and the pressure term was absent in the Galerkin approximations.

The objective of this paper is to define semidiscrete finite element approximations, prove the convergence of finite element solutions, and derive error estimates for the finite element approximations. We point out that finite element basis functions in general are not divergence-free, and finite element formulations must be studied with the pressure term. The proof for the convergence of finite element solutions provides an alternative proof to that found in [10] for the existence of a weak solution; the results of this paper do not rely on those of [10] concerning the existence of a divergence-free weak solution. Moreover, the regularity and compatibility assumptions made on the data in this paper lead to a stronger solution. The details for the divergence-free Galerkin approximations of [10] and the discretely divergence-free finite element approximations are sufficiently different so that separate treatments are warranted.

A few technical aspects contained in this paper are particularly noteworthy: the finite element initial conditions are defined asymmetrically about the two subdomains Ω_1 and Ω_2 ; two inf-sup conditions are verified that facilitate the analysis of certain steady-state saddle point problems (these inf-sup conditions are also useful in dealing with approximations of mixed boundary value problems for the Stokes equations); and error estimates for a weighted L^2 projection onto discretely divergence-free spaces are derived.

The plan of the paper is as follows. In section 2, we recall relevant results of [10], in particular the weak formulations and the existence theorems. In section 3, we define semidiscrete finite element approximations and establish the existence of and a priori estimates for the finite element solutions. In section 4, we show the convergence of finite element solutions and derive error estimates.

2. Notations and results concerning divergence-free weak formulations.

In this section we will recall the notation, weak formulations, and existence results of [10].

Throughout this paper, C denotes a positive constant, depending on the domains Ω , Ω_1 , and Ω_2 , whose meaning and value changes with context. $H^s(\mathcal{D})$, $s \in \mathbb{R}$, denotes the standard Sobolev space of order s with respect to the set \mathcal{D} equipped with the standard norm $\|\cdot\|_{s,\mathcal{D}}$. Vector-valued Sobolev spaces are denoted by $\mathbf{H}^s(\mathcal{D})$, with norms still denoted by $\|\cdot\|_{s,\mathcal{D}}$. $H_0^1(\mathcal{D})$ denotes the space of functions belonging to $H^1(\mathcal{D})$ that vanish on the boundary $\partial\mathcal{D}$ of \mathcal{D} ; $\mathbf{H}_0^1(\mathcal{D})$ denotes the vector-valued counterpart.

We will use the following L^2 inner product notations on scalar and vector-valued L^2 spaces:

$$[p, q]_{\mathcal{D}} = \int_{\mathcal{D}} pq \, d\mathcal{D} \quad \forall p, q \in L^2(\mathcal{D}), \quad [\mathbf{u}, \mathbf{v}]_{\mathcal{D}} = \int_{\mathcal{D}} \mathbf{u} \cdot \mathbf{v} \, d\mathcal{D} \quad \forall \mathbf{u}, \mathbf{v} \in \mathbf{L}^2(\mathcal{D}),$$

where the spatial set \mathcal{D} is Ω or Γ_0 or Ω_i , for $i = 1, 2$.

We introduce the function spaces

$$X_i = [\mathbf{H}_0^1(\Omega)]|_{\Omega_i} \quad \text{with the norm } \|\cdot\|_{X_i} = \|\cdot\|_{1,\Omega_i}, \quad i = 1, 2,$$

and

$$\Psi = \{\boldsymbol{\eta} \in \mathbf{H}_0^1(\Omega) : \operatorname{div} \boldsymbol{\eta} = 0 \text{ in } \Omega_1\} \quad \text{with the norm } \|\cdot\|_{1,\Omega}.$$

We define the weighted $\mathbf{L}^2(\Omega)$ inner product $[[\cdot, \cdot]]$ by

$$(2.1) \quad [[\boldsymbol{\xi}, \boldsymbol{\eta}]] = [\rho_1 \boldsymbol{\xi}, \boldsymbol{\eta}]_{\Omega_1} + [\rho_2 \boldsymbol{\xi}, \boldsymbol{\eta}]_{\Omega_2} \quad \forall \boldsymbol{\xi}, \boldsymbol{\eta} \in \mathbf{L}^2(\Omega).$$

We denote by $\langle\langle \cdot, \cdot \rangle\rangle$ the duality pairing between Ψ^* and Ψ that is generated from the weighted $\mathbf{L}^2(\Omega)$ inner product $[[\cdot, \cdot]]$. The norm on the dual space Ψ^* is defined in the conventional manner:

$$\|\mathbf{g}\|_{\Psi^*} = \sup_{\boldsymbol{\eta} \in \Psi, \|\boldsymbol{\eta}\|_{1,\Omega} \leq 1} |\langle\langle \mathbf{g}, \boldsymbol{\eta} \rangle\rangle| \quad \forall \mathbf{g} \in \Psi^*.$$

We define the bilinear forms

$$\begin{aligned} a_1[\mathbf{u}, \mathbf{v}] &= 2 \int_{\Omega_1} \mu_1 (\nabla \mathbf{u} + \nabla \mathbf{u}^T) : (\nabla \mathbf{v} + \nabla \mathbf{v}^T) d\Omega \quad \forall \mathbf{u}, \mathbf{v} \in X_1, \\ a_2[\mathbf{u}, \mathbf{v}] &= \int_{\Omega_2} \left\{ 2\mu_2 (\nabla \mathbf{u} + \nabla \mathbf{u}^T) : (\nabla \mathbf{v} + \nabla \mathbf{v}^T) + \lambda_2 (\nabla \cdot \mathbf{u})(\nabla \cdot \mathbf{v}) \right\} d\Omega \quad \forall \mathbf{u}, \mathbf{v} \in X_2, \\ b[\mathbf{v}, q] &= - \int_{\Omega_1} q \nabla \cdot \mathbf{v} d\Omega \quad \forall \mathbf{v} \in X_1, \forall q \in L^2(\Omega_1). \end{aligned}$$

It can be verified with the help of Korn's inequalities [31, pp. 31, 120] that for $i = 1, 2$,

$$(2.2) \quad a_i[\boldsymbol{\eta}, \boldsymbol{\eta}] \geq k_i \|\boldsymbol{\eta}\|_{1, \Omega_i}^2 \quad \forall \boldsymbol{\eta} \in X_i \quad \text{if } \text{meas}(\Gamma_i) \neq 0$$

and

$$(2.3) \quad [\boldsymbol{\eta}, \boldsymbol{\eta}]_{\Omega_i} + a_i[\boldsymbol{\eta}, \boldsymbol{\eta}] \geq k_i \|\boldsymbol{\eta}\|_{1, \Omega_i}^2 \quad \forall \boldsymbol{\eta} \in X_i \quad \text{if } \text{meas}(\Gamma_i) = 0.$$

The bounded bilinear form $b[\cdot, \cdot]$ was shown in [10] to satisfy the inf-sup conditions

$$(2.4) \quad \inf_{q \in L^2(\Omega_1)} \sup_{\boldsymbol{\eta} \in \mathbf{H}_0^1(\Omega)} \frac{b[\boldsymbol{\eta}, q]}{\|\boldsymbol{\eta}\|_{1, \Omega} \|q\|_{0, \Omega_1}} \geq k_b$$

and

$$(2.5) \quad \inf_{q \in L^2(\Omega_1)} \sup_{\mathbf{v} \in X_1} \frac{b[\mathbf{v}, q]}{\|\mathbf{v}\|_{1, \Omega_1} \|q\|_{0, \Omega_1}} \geq k_b,$$

where $k_b > 0$ is a constant.

For functions that also depend on time, we introduce the space $L^2(0, T; X)$ that consists of L^2 -integrable functions from $[0, T]$ into the space X and which is equipped with the norm

$$\left(\int_0^t \|f\|_X^2 dt \right)^{1/2}.$$

Similarly, we introduce the space $C(0, T; X)$ that consists of continuous functions from $[0, T]$ into the space X and which is equipped with the norm

$$\sup_{t \in [0, T]} \|f\|_X.$$

The divergence-free weak formulation for (1.1)–(1.4) was defined in [10] as follows. Given

$$(2.6) \quad \begin{cases} \mathbf{f}_1 \in C([0, T]; \mathbf{L}^2(\Omega_1)), & \mathbf{f}_2 \in C([0, T]; \mathbf{L}^2(\Omega_2)), & \mathbf{u}_0 \in X_2, \\ \mathbf{v}_0 \in X_1, & \text{div } \mathbf{v}_0 = 0 \text{ in } \Omega_1, & \mathbf{u}_1 \in X_2, & \mathbf{v}_0|_{\Gamma_0} = \mathbf{u}_1|_{\Gamma_0}, \end{cases}$$

seek a pair (\mathbf{v}, \mathbf{u}) such that

$$(2.7) \quad (\mathbf{v}, \mathbf{u}) \in L^2(0, T; X_1) \times L^2(0, T; X_2), \quad \text{div } \mathbf{v} = 0,$$

$$(2.8) \quad \begin{aligned} & \frac{d}{dt} \left(\rho_1 [\mathbf{v}, \boldsymbol{\eta}]_{\Omega_1} + \rho_2 [\partial_t \mathbf{u}, \boldsymbol{\eta}]_{\Omega_2} \right) + a_1[\mathbf{v}, \boldsymbol{\eta}] + a_2[\mathbf{u}, \boldsymbol{\eta}] \\ & = \rho_1 [\mathbf{f}_1, \boldsymbol{\eta}]_{\Omega_1} + \rho_2 [\mathbf{f}_2, \boldsymbol{\eta}]_{\Omega_2} \quad \forall \boldsymbol{\eta} \in \boldsymbol{\Psi}, \end{aligned}$$

$$(2.9) \quad \mathbf{v}|_{t=0} = \mathbf{v}_0, \quad \mathbf{u}|_{t=0} = \mathbf{u}_0, \quad \mathbf{u}_t|_{t=0} = \mathbf{u}_1,$$

and

$$(2.10) \quad \int_0^t \mathbf{v}(s)|_{\Gamma_0} ds = \mathbf{u}(t)|_{\Gamma_0} - \mathbf{u}_0|_{\Gamma_0} \quad \text{a.e. } t.$$

The “natural” interface condition (1.4) is built into (2.8), and the “essential” interface condition (1.3) is enforced weakly in the sense of (2.10).

By defining

$$(2.11) \quad \boldsymbol{\xi} = \begin{cases} \mathbf{v} & \text{in } \Omega_1, \\ \mathbf{u}_t & \text{in } \Omega_2, \end{cases} \quad \boldsymbol{\xi}_0 = \begin{cases} \mathbf{v}_0 & \text{in } \Omega_1, \\ \mathbf{u}_1 & \text{in } \Omega_2, \end{cases} \quad \text{and} \quad \mathbf{f} = \begin{cases} \mathbf{f}_1 & \text{in } \Omega_1, \\ \mathbf{f}_2 & \text{in } \Omega_2, \end{cases}$$

(2.7)–(2.10) was conveniently recast in [10] into the following equivalent, auxiliary divergence-free weak formulation: seek a $\boldsymbol{\xi}$ such that

$$(2.12) \quad \begin{aligned} \boldsymbol{\xi} &\in L^2(0, T; \mathbf{L}^2(\Omega)), & \partial_t \boldsymbol{\xi} &\in L^2(0, T; \boldsymbol{\Psi}^*), \\ \boldsymbol{\xi}|_{\Omega_1} &\in L^2(0, T; X_1), & \operatorname{div} \boldsymbol{\xi}|_{\Omega_1} &= 0, & \int_0^t \boldsymbol{\xi}(s)|_{\Omega_2} ds &\in L^2(0, T; X_2), \end{aligned}$$

(2.13)

$$\langle \langle \boldsymbol{\xi}_t, \boldsymbol{\eta} \rangle \rangle + a_1[\boldsymbol{\xi}, \boldsymbol{\eta}] + a_2 \left[\int_0^t \boldsymbol{\xi}(s) ds, \boldsymbol{\eta} \right] = [[\mathbf{f}, \boldsymbol{\eta}]] - a_2[\mathbf{u}_0, \boldsymbol{\eta}] \quad \forall \boldsymbol{\eta} \in \boldsymbol{\Psi}, \quad \text{a.e. } t,$$

$$(2.14) \quad \boldsymbol{\xi}(0) = \boldsymbol{\xi}_0,$$

and

$$(2.15) \quad \int_0^t (\boldsymbol{\xi}(s)|_{\Omega_1})|_{\Gamma_0} ds = \int_0^t (\boldsymbol{\xi}(s)|_{\Omega_2})|_{\Gamma_0} ds \quad \text{a.e. } t.$$

The existence and uniqueness of a solution for the auxiliary problem (2.12)–(2.15) was proved in [10].

THEOREM 2.1. *Assume that $\mathbf{f}_1, \mathbf{v}_0, \mathbf{f}_2$, and \mathbf{u}_0 satisfy (2.6). Then, there exists a unique solution $\boldsymbol{\xi}$ for (2.12)–(2.15). Moreover, $\boldsymbol{\xi}$ satisfies the estimates*

$$(2.16) \quad \begin{aligned} &\|\boldsymbol{\xi}(t)\|_{0, \Omega}^2 + \|\boldsymbol{\xi}\|_{L^2(0, T; \mathbf{H}^1(\Omega_1))}^2 + \left\| \int_0^t \boldsymbol{\xi}(s) ds \right\|_{\mathbf{H}^1(\Omega_2)}^2 \\ &\leq C e^{CT} (\|\mathbf{f}\|_{L^2(0, T; \mathbf{L}^2(\Omega))}^2 + \|\mathbf{u}_0\|_{1, \Omega_2}^2 + \|\mathbf{v}_0\|_{1, \Omega_1}^2 + \|\mathbf{u}_1\|_{1, \Omega_2}^2) \quad \forall t \in [0, T] \end{aligned}$$

and

$$(2.17) \quad \begin{aligned} &\|\partial_t \boldsymbol{\xi}\|_{L^2(0, T; \boldsymbol{\Psi}^*)}^2 \\ &\leq C e^{CT} (\|\mathbf{f}\|_{L^2(0, T; \mathbf{L}^2(\Omega))}^2 + \|\mathbf{u}_0\|_{1, \Omega_2}^2 + \|\mathbf{v}_0\|_{1, \Omega_1}^2 + \|\mathbf{u}_1\|_{1, \Omega_2}^2). \end{aligned}$$

Using relation (2.11) reversely, i.e., setting $\mathbf{v} = \boldsymbol{\xi}|_{\Omega_1}$ and $\mathbf{u} = \mathbf{u}_0 + \int_0^t \boldsymbol{\xi}(s)|_{\Omega_2} ds$, Theorem 2.1 immediately yields the following existence result for (2.7)–(2.10).

THEOREM 2.2. *Assume that $\mathbf{f}_1, \mathbf{v}_0, \mathbf{f}_2, \mathbf{u}_0$, and \mathbf{u}_1 satisfy (2.6). Then, there exists a unique solution $(\mathbf{v}, \mathbf{u}) \in L^2(0, T; X_1) \times L^2(0, T; X_2)$ for (2.7)–(2.10), where (2.8) holds in the sense of distributions on $(0, T)$. Moreover,*

$$(2.18) \quad \begin{aligned} & \|\mathbf{v}(t)\|_{0, \Omega_1}^2 + \|\mathbf{u}_t(t)\|_{0, \Omega_2}^2 + \|\mathbf{v}\|_{L^2(0, T; \mathbf{H}^1(\Omega_1))}^2 + \|\mathbf{u}(t)\|_{\mathbf{H}^1(\Omega_2)}^2 \\ & \leq C e^{CT} (\|\mathbf{f}\|_{L^2(0, T; \mathbf{L}^2(\Omega))}^2 + \|\mathbf{u}_0\|_{1, \Omega_2}^2 + \|\mathbf{v}_0\|_{0, \Omega_1}^2 + \|\mathbf{u}_1\|_{0, \Omega_2}^2) \quad \forall t \in [0, T]. \end{aligned}$$

The existence of a stronger solution and an L^2 -integrable pressure was also established in [10].

THEOREM 2.3. *Assume that $\mathbf{f}_1, \mathbf{v}_0, \mathbf{f}_2, \mathbf{u}_0$, and \mathbf{u}_1 satisfy (2.6) and*

$$\partial_t \mathbf{f}_i \in L^2(0, T; \mathbf{L}^2(\Omega_i)), \quad i = 1, 2, \quad \mathbf{v}_0 \in \mathbf{H}^2(\Omega_1), \quad \mathbf{u}_1 \in \mathbf{H}^1(\Omega_2), \quad \mathbf{u}_0 \in \mathbf{H}^2(\Omega_2).$$

Assume further that there exists a $p_0 \in H^1(\Omega_1)$ such that

$$(p_0 \mathbf{n}_1 - \mu_1 (\nabla \mathbf{v}_0 + \nabla \mathbf{v}_0^T) \cdot \mathbf{n}_1)|_{\Gamma_0} = (\mu_2 (\nabla \mathbf{u}_0 + \nabla \mathbf{u}_0^T) \cdot \mathbf{n}_2 + (\lambda_2 + \mu_2) (\operatorname{div} \mathbf{u}_0) \mathbf{n}_2)|_{\Gamma_0},$$

where \mathbf{n}_i denotes the outward-pointing normal along $\partial \Omega_i$. Then, the solution (\mathbf{v}, \mathbf{u}) to (2.7)–(2.10) satisfies

$$\mathbf{v} \in L^\infty(0, T; \mathbf{L}^2(\Omega_1)) \cap L^2(0, T; X_1), \quad \mathbf{u} \in L^\infty(0, T; X_2),$$

$$\mathbf{v}_t \in L^\infty(0, T; \mathbf{L}^2(\Omega_1)) \cap L^2(0, T; X_1), \quad \mathbf{u}_t \in L^\infty(0, T; X_2), \quad \mathbf{u}_{tt} \in L^\infty(0, T; \mathbf{L}^2(\Omega_2)),$$

and

$$\begin{aligned} & \|\partial_t \mathbf{v}(t)\|_{0, \Omega_1}^2 + \|\partial_{tt} \mathbf{u}(t)\|_{0, \Omega_2}^2 + \|\partial_t \mathbf{v}\|_{L^2(0, T; X_1)}^2 + \|\partial_t \mathbf{u}(t)\|_{1, \Omega_2}^2 \\ & \leq C e^{CT} (\|\mathbf{f}\|_{H^1(0, T; \mathbf{L}^2(\Omega))}^2 + \|\mathbf{u}_0\|_{2, \Omega_2}^2 + \|\mathbf{v}_0\|_{2, \Omega_1}^2 + \|p_0\|_{1, \Omega_1}^2 + \|\mathbf{u}_1\|_{1, \Omega_2}^2) \quad \forall t \in [0, T]. \end{aligned}$$

Furthermore, there exists a unique $p \in L^2(0, T; L^2(\Omega_1))$ such that

$$(2.19) \quad \begin{aligned} & \rho_1 [\mathbf{v}_t, \boldsymbol{\eta}]_{\Omega_1} + b[\boldsymbol{\eta}, p] + a_1[\mathbf{v}, \boldsymbol{\eta}] + \rho_2 [\mathbf{u}_{tt}, \boldsymbol{\eta}]_{\Omega_2} + a_2[\mathbf{u}, \boldsymbol{\eta}] \\ & = \rho_1 [\mathbf{f}_1, \boldsymbol{\eta}]_{\Omega_1} + \rho_2 [\mathbf{f}_2, \boldsymbol{\eta}]_{\Omega_2} \quad \forall \boldsymbol{\eta} \in \mathbf{H}_0^1(\Omega), \quad a.e. \ t \end{aligned}$$

and

$$\|p\|_{L^2(0, T; L^2(\Omega_1))} \leq C e^{CT} (\|\mathbf{f}\|_{H^1(0, T; \mathbf{L}^2(\Omega))}^2 + \|\mathbf{u}_0\|_{2, \Omega_2}^2 + \|\mathbf{v}_0\|_{2, \Omega_1}^2 + \|p_0\|_{1, \Omega_1}^2 + \|\mathbf{u}_1\|_{1, \Omega_2}^2).$$

3. Semidiscrete finite element approximations. In this section we will define semidiscrete finite element approximations, prove the existence of finite element solutions on discretely divergence-free spaces and derive energy estimates, and establish the existence of a discrete pressure by verifying inf-sup conditions for finite element space pairs.

As alluded to previously, finite element solutions in general are not divergence-free, and finite element formulations should include the pressure term. Of course, the corresponding continuous weak formulation should also contain the pressure term. Such a weak formulation requires additional regularity on \mathbf{v}_t and \mathbf{u}_{tt} . The continuous weak formulation we consider is as follows: given $\mathbf{f}_1, \mathbf{v}_0, \mathbf{f}_2$, and \mathbf{u}_0 satisfying (2.6), seek a triplet $(\mathbf{v}, p, \mathbf{u})$ such that

$$(3.1) \quad (\mathbf{v}, p, \mathbf{u}) \in L^2(0, T; X_1) \times L^2(0, T; L^2(\Omega_1)) \times L^2(0, T; X_2),$$

$$(3.2) \quad \mathbf{v}_t \in L^2(0, T; \mathbf{L}^2(\Omega_1)), \quad \mathbf{u}_t \in L^2(0, T; X_2), \quad \mathbf{u}_{tt} \in L^2(0, T; \mathbf{L}^2(\Omega_2)),$$

$$(3.3) \quad \begin{aligned} & \rho_1[\mathbf{v}_t, \boldsymbol{\eta}]_{\Omega_1} + b[\boldsymbol{\eta}, p] + a_1[\mathbf{v}, \boldsymbol{\eta}] + \rho_2[\mathbf{u}_{tt}, \boldsymbol{\eta}]_{\Omega_2} + a_2[\mathbf{u}, \boldsymbol{\eta}] \\ & = \rho_1[\mathbf{f}_1, \boldsymbol{\eta}]_{\Omega_1} + \rho_2[\mathbf{f}_2, \boldsymbol{\eta}]_{\Omega_2} \quad \forall \boldsymbol{\eta} \in \mathbf{H}_0^1(\Omega), \text{ a.e. } t \in [0, T], \end{aligned}$$

$$(3.4) \quad b[\mathbf{v}, q] = 0 \quad \forall q \in L^2(\Omega_1), \text{ a.e. } t \in [0, T],$$

$$(3.5) \quad \mathbf{v}|_{t=0} = \mathbf{v}_0, \quad \mathbf{u}|_{t=0} = \mathbf{u}_0, \quad \mathbf{u}_t|_{t=0} = \mathbf{u}_1,$$

$$(3.6) \quad \mathbf{v}|_{\Gamma_0} = \mathbf{u}_t|_{\Gamma_0} \quad \text{a.e. } t.$$

We will define finite element approximations to (3.3)–(3.6). By showing the convergence of finite element solutions, we establish the existence of a solution for (3.1)–(3.6). For reasons connected with the derivation of the regularity results (3.2), we will define finite element initial conditions in a nonstandard manner.

3.1. Finite element discretization. In what follows we assume that Ω_1 and Ω_2 are two-dimensional polygons or three-dimensional polyhedra. Let h denote a discretization parameter associated with the triangulation $\mathcal{T}^h(\Omega)$ of Ω . We assume that elements of \mathcal{T}^h do not cross the interface Γ_0 . We assume that the triangulation \mathcal{T}^h consists of triangular elements in two dimensions or tetrahedral elements in three dimensions, though our results can be extended to other types of triangulations. Furthermore, we assume that there exists a triangulation $\mathcal{T}^{h_0}(\Omega)$ such that, for each $h < h_0$, $\mathcal{T}^h(\Omega)$ is a refinement of $\mathcal{T}^{h_0}(\Omega)$.

For each h , we choose $X^h \subset \mathbf{C}(\bar{\Omega}) \cap \mathbf{H}_0^1(\Omega)$ and $Q_1^h \subset L^2(\Omega_1)$ as finite element subspaces over the triangulation $\mathcal{T}^h(\Omega)$. We assume that X^h contains piecewise linear functions. We set

$$X_i^h = X^h|_{\Omega_i}, \quad i = 1, 2,$$

and

$$\boldsymbol{\Psi}^h = \{\boldsymbol{\eta}_h \in X^h : b[\boldsymbol{\eta}_h, q_h] = 0 \forall q_h \in Q_1^h\}.$$

We assume that the finite element spaces X_1^h , X_2^h , and Q_1^h satisfy the standard approximation properties [5]; i.e., there exist an integer $k > 0$ and constant $C > 0$ such that

$$(3.7) \quad \inf_{\mathbf{v}^h \in X_i^h} \|\mathbf{v} - \mathbf{v}^h\|_{0, \Omega_i} \leq Ch^{r+1} \|\mathbf{v}\|_{r+1, \Omega_i} \quad \forall \mathbf{v} \in \mathbf{H}^{r+1}(\Omega_i) \cap X_i, \quad r \in [0, k],$$

$$(3.8) \quad \inf_{\mathbf{v}^h \in X_i^h} \|\mathbf{v} - \mathbf{v}^h\|_{1, \Omega_i} \leq Ch^r \|\mathbf{v}\|_{r+1, \Omega_i} \quad \forall \mathbf{v} \in \mathbf{H}^{r+1}(\Omega_i) \cap X_i, \quad r \in [0, k],$$

and

$$(3.9) \quad \inf_{q^h \in Q_1^h} \|q - q^h\|_{0, \Omega_1} \leq Ch^r \|p\|_{r, \Omega_1} \quad \forall q \in H^r(\Omega_1), \quad r \in [0, k].$$

Also, X^h satisfies the approximation properties

$$(3.10) \quad \inf_{\boldsymbol{\eta}^h \in X^h} \|\boldsymbol{\eta} - \boldsymbol{\eta}^h\|_{0,\Omega} \leq Ch^{r+1} \|\boldsymbol{\eta}\|_{r+1,\Omega} \quad \forall \boldsymbol{\eta} \in \mathbf{H}^{r+1}(\Omega) \cap \mathbf{L}^2(\Omega), \quad r \in [0, k],$$

and

$$(3.11) \quad \inf_{\boldsymbol{\eta}^h \in X^h} \|\boldsymbol{\eta} - \boldsymbol{\eta}^h\|_{1,\Omega_i} \leq Ch^r \|\boldsymbol{\eta}\|_{r+1,\Omega} \quad \forall \boldsymbol{\eta} \in \mathbf{H}^{r+1}(\Omega) \cap \mathbf{H}_0^1(\Omega), \quad r \in [0, k].$$

We assume that the finite element pair $\{\tilde{X}_1^h, M^h\} \equiv \{X_1^h \cap \mathbf{H}_0^1(\Omega_1), Q_1^h \cap L_0^2(\Omega_1)\}$ satisfies the discrete inf-sup condition

$$(3.12) \quad \inf_{\mathbf{v}^h \in M^h(\Omega_1)} \sup_{q^h \in \tilde{X}_1^h(\Omega_1)} \frac{b[\mathbf{v}^h, q^h]}{\|\mathbf{v}^h\|_{1,\Omega_1} \|q^h\|_{0,\Omega_1}} \geq C.$$

Choices of finite element spaces satisfying (3.12) are well known [19]. Note that functions in \tilde{X}_1^h vanish on Γ_0 .

We also assume that triangulations are uniformly regular so that the following inverse inequalities hold:

$$(3.13) \quad \begin{aligned} \|\mathbf{v}^h\|_{1,\Omega} &\leq Ch^{-1} \|\mathbf{v}^h\|_{0,\Omega} \quad \forall \mathbf{v}^h \in X^h; \\ \|\mathbf{v}^h\|_{1,\Omega_i} &\leq Ch^{-1} \|\mathbf{v}^h\|_{0,\Omega_i} \quad \forall \mathbf{v}^h \in X_i^h, \quad i = 1, 2. \end{aligned}$$

Semidiscrete finite element approximations of the weak form (3.3)–(3.6) are defined as follows: seek $(\mathbf{v}^h, p^h, \mathbf{u}^h) \in C^1([0, T]; X_1^h) \times C([0, T]; Q_1^h) \times C^1([0, T]; X_2^h)$ such that

$$(3.14) \quad \begin{aligned} \rho_1[\partial_t \mathbf{v}_h, \boldsymbol{\eta}_h]_{\Omega_1} + b[\boldsymbol{\eta}_h, p_h] + a_1[\mathbf{v}_h, \boldsymbol{\eta}_h] + \rho_2[\partial_{tt} \mathbf{u}_h, \boldsymbol{\eta}_h]_{\Omega_2} + a_2[\mathbf{u}_h, \boldsymbol{\eta}_h] \\ = \rho_1[\mathbf{f}_1, \boldsymbol{\eta}_h]_{\Omega_1} + \rho_2[\mathbf{f}_2, \boldsymbol{\eta}_h]_{\Omega_2} \quad \forall \boldsymbol{\eta}_h \in X^h, \quad \text{a.e. } t, \end{aligned}$$

$$(3.15) \quad b[\mathbf{v}_h, q_h] = 0 \quad \forall q_h \in Q_1^h, \quad \text{a.e. } t,$$

$$(3.16) \quad \mathbf{v}_h|_{\Gamma_0} = \partial_t \mathbf{u}_h|_{\Gamma_0} \quad \text{a.e. } t \in [0, T],$$

$$(3.17) \quad \mathbf{v}_h|_{t=0} = \mathbf{v}_{0,h}, \quad \mathbf{u}_h|_{t=0} = \mathbf{u}_{0,h}, \quad \partial_t \mathbf{u}_h|_{t=0} = \mathbf{u}_{1,h},$$

where $\mathbf{v}_{0,h} \in \boldsymbol{\Psi}^h|_{\Omega_1}$, $\mathbf{u}_{0,h} \in X_2^h$, and $\mathbf{u}_{1,h} \in X_2^h$ are finite element approximations of \mathbf{v}_0 , \mathbf{u}_0 , and \mathbf{u}_1 , respectively. We assume that $(\mathbf{v}_{0,h}, \mathbf{u}_{1,h})$ satisfies

$$(3.18) \quad b[\mathbf{v}_{0,h}, q_h] = 0 \quad \forall q_h \in Q_1^h, \quad \mathbf{v}_{0,h}|_{\Gamma_0} = \mathbf{u}_{1,h}|_{\Gamma_0}$$

and that $\mathbf{u}_{0,h}$ is defined by

$$(3.19) \quad a_2[\mathbf{u}_{0,h}, \mathbf{w}_h] = a_2[\mathbf{u}_0, \mathbf{w}_h] \quad \forall \mathbf{w}_h \in X_2^h.$$

3.2. The existence of discretely divergence-free finite element solutions.

The existence of finite element solutions $\{(\mathbf{v}^h, \mathbf{u}^h)\}$ can be established in a manner analogous to the analysis of the Galerkin approximations $\{(\mathbf{v}_m, \mathbf{u}_m)\}$ in [10]. However, it should be noted that finite element approximations are not special cases of the Galerkin approximations due to the fact that the basis functions used in the Galerkin approximations are divergence-free in Ω_1 , whereas the finite element solutions are only discretely divergence-free in Ω_1 in the sense of (3.15), i.e., they belong to the space of discretely divergence-free functions Ψ^h .

We first formulate auxiliary semidiscrete finite element approximations on the discretely divergence-free space Ψ^h . Through the relation

$$(3.20) \quad \boldsymbol{\xi}_h = \begin{cases} \mathbf{v}_h & \text{in } \Omega_1, \\ \partial_t \mathbf{u}_h & \text{in } \Omega_2, \end{cases}$$

we see that (3.14)–(3.19) can be recast into the system

$$(3.21) \quad \begin{aligned} & \rho_1[\partial_t \boldsymbol{\xi}_h, \boldsymbol{\eta}_h]_{\Omega_1} + \rho_2[\partial_t \boldsymbol{\xi}_h, \boldsymbol{\eta}_h]_{\Omega_2} + a_1[\boldsymbol{\xi}_h, \boldsymbol{\eta}_h] + a_2 \left[\int_0^t \boldsymbol{\xi}_h(s) ds, \boldsymbol{\eta}_h \right] \\ & = \rho_1[\mathbf{f}_1, \boldsymbol{\eta}_h]_{\Omega_1} + \rho_2[\mathbf{f}_2, \boldsymbol{\eta}_h]_{\Omega_2} - a_2[\mathbf{u}_0, \boldsymbol{\eta}_h] \quad \forall \boldsymbol{\eta}_h \in \Psi^h, t \in [0, T] \end{aligned}$$

and

$$(3.22) \quad \boldsymbol{\xi}_h(0) = \boldsymbol{\xi}_{0,h} \equiv \begin{cases} \mathbf{v}_{0,h} & \text{in } \Omega_1, \\ \mathbf{u}_{1,h} & \text{in } \Omega_2. \end{cases}$$

Let $\{\boldsymbol{\psi}_j^h\}_{j=1}^{J_h}$ be a finite element basis for Ψ^h . Assumption (3.18) implies that $\boldsymbol{\xi}_{0,h} \in \Psi^h$, so that we can write

$$\boldsymbol{\xi}_{0,h} = \sum_{j=1}^{J_h} d_j \boldsymbol{\psi}_j^h.$$

The solution $\boldsymbol{\xi}^h \in C([0, T]; \Psi^h)$ for (3.21)–(3.22) can be expressed in the form

$$(3.23) \quad \boldsymbol{\xi}_h = \sum_{j=1}^{J_h} g_j^h(t) \boldsymbol{\psi}_j^h(\mathbf{x})$$

so that system (3.21)–(3.22) is equivalent to the following linear system of ordinary differential equations for $\{g_j^h\}_{j=1}^{J_h}$:

$$\begin{cases} \sum_{j=1}^{J_h} [[\boldsymbol{\psi}_j^h, \boldsymbol{\psi}_i^h]] \frac{d}{dt} g_j^h(t) + \sum_{j=1}^{J_h} a_1[\boldsymbol{\psi}_j^h, \boldsymbol{\psi}_i^h] g_j^h(t) + \sum_{j=1}^{J_h} a_2[\boldsymbol{\psi}_j^h, \boldsymbol{\psi}_i^h] \int_0^t g_j^h(s) ds \\ = [\rho_1 \mathbf{f}_1(t), \boldsymbol{\psi}_i^h]_{\Omega_1} + [\rho_2 \mathbf{f}_2(t), \boldsymbol{\psi}_i^h]_{\Omega_2} - a_2[\mathbf{u}_0, \boldsymbol{\psi}_i^h], & i = 1, \dots, J_h, t \in [0, T], \\ g_i^h(0) = d_i, & i = 1, \dots, J_h. \end{cases}$$

We have the following results concerning the existence of and a priori estimates for a finite element solution $\boldsymbol{\xi}_h$ of (3.21)–(3.22). The proof is the same as that in [10] for the Galerkin approximations and thus is omitted.

THEOREM 3.1. *Assume that $\mathbf{f}_1, \mathbf{v}_0, \mathbf{f}_2, \mathbf{u}_0$, and \mathbf{u}_1 satisfy (2.6). Then, there exists a unique function $\boldsymbol{\xi}_h \in C^1([0, T]; \boldsymbol{\Psi}^h)$ which satisfies (3.21)–(3.22) and the estimate*

$$(3.24) \quad \begin{aligned} & \|\boldsymbol{\xi}_h(t)\|_{0,\Omega}^2 + \|\boldsymbol{\xi}_h\|_{L^2(0,T;\mathbf{H}^1(\Omega_1))}^2 + \left\| \int_0^t \boldsymbol{\xi}_h(s) ds \right\|_{\mathbf{H}^1(\Omega_2)}^2 \\ & \leq C e^{CT} (\|\mathbf{f}\|_{L^2(0,T;L^2(\Omega))}^2 + \|\mathbf{u}_0\|_{1,\Omega_2}^2 + \|\mathbf{v}_{0,h}\|_{0,\Omega_1}^2 + \|\mathbf{u}_{1,h}\|_{0,\Omega_2}^2) \forall t \in [0, T]. \end{aligned}$$

Setting $\mathbf{v}_h = \boldsymbol{\xi}_h|_{\Omega_1}$, $\mathbf{u}_h = \mathbf{u}_{0,h} + \int_0^t \boldsymbol{\xi}_h(s)|_{\Omega_2} ds$ and using (3.19), we immediately obtain the existence of a $(\mathbf{v}_h, \mathbf{u}_h)$ satisfying the discretely divergence-free version of (3.14)–(3.19), as follows.

THEOREM 3.2. *Assume that $\mathbf{f}_1, \mathbf{v}_0, \mathbf{f}_2, \mathbf{u}_0$, and \mathbf{u}_1 satisfy (2.6). Then, there exists a unique $(\mathbf{v}_h, \mathbf{u}_h) \in C^1([0, T]; \boldsymbol{\Psi}^h|_{\Omega_1}) \times C^2([0, T]; X_2)$ satisfying*

$$(3.25) \quad \begin{aligned} & \rho_1[\partial_t \mathbf{v}_h, \boldsymbol{\eta}_h]_{\Omega_1} + a_1[\mathbf{v}, \boldsymbol{\eta}_h] + \rho_2[\partial_{tt} \mathbf{u}_h, \boldsymbol{\eta}_h]_{\Omega_2} + a_2[\mathbf{u}_h, \boldsymbol{\eta}_h] \\ & = \rho_1[\mathbf{f}_1, \boldsymbol{\eta}_h]_{\Omega_1} + \rho_2[\mathbf{f}_2, \boldsymbol{\eta}_h]_{\Omega_2} \quad \forall \boldsymbol{\eta}_h \in \boldsymbol{\Psi}^h, t \in [0, T] \end{aligned}$$

and (3.15)–(3.19). Moreover, the following estimate holds:

$$(3.26) \quad \begin{aligned} & \|\mathbf{v}_h(t)\|_{0,\Omega_1}^2 + \|\partial_t \mathbf{u}_h(t)\|_{0,\Omega_2}^2 + \|\mathbf{v}_h\|_{L^2(0,T;\mathbf{H}^1(\Omega_1))}^2 + \|\mathbf{u}_h\|_{\mathbf{H}^1(\Omega_2)}^2 \\ & \leq C e^{CT} (\|\mathbf{f}\|_{L^2(0,T;L^2(\Omega))}^2 + \|\mathbf{u}_0\|_{1,\Omega_2}^2 + \|\mathbf{v}_{0,h}\|_{0,\Omega_1}^2 + \|\mathbf{u}_{1,h}\|_{0,\Omega_2}^2) \forall t \in [0, T]. \end{aligned}$$

3.3. The discrete inf-sup conditions and discrete pressure fields. We have proved the existence of a finite element solution in the discretely divergence-free formulation consisting of (3.25) and (3.15)–(3.19). We will show the existence of a discrete pressure p_h such that (3.14) holds. A crucial step towards this goal is the verification of discrete inf-sup conditions. The discrete inf-sup conditions will also play a role in deriving strong energy estimates in a subsequent section.

We rewrite (3.14) as

$$(3.27) \quad \begin{aligned} b[\boldsymbol{\eta}_h, p_h] & = -\rho_1[\partial_t \mathbf{v}_h, \boldsymbol{\eta}_h]_{\Omega_1} - a_1[\mathbf{v}, \boldsymbol{\eta}_h] - \rho_2[\partial_{tt} \mathbf{u}_h, \boldsymbol{\eta}_h]_{\Omega_2} - a_2[\mathbf{u}_h, \boldsymbol{\eta}_h] \\ & + \rho_1[\mathbf{f}_1, \boldsymbol{\eta}_h]_{\Omega_1} + \rho_2[\mathbf{f}_2, \boldsymbol{\eta}_h]_{\Omega_2} \quad \forall \boldsymbol{\eta}_h \in X^h, t \in [0, T]. \end{aligned}$$

In terms of the auxiliary variable $\boldsymbol{\xi}_h$, (3.27) is equivalent to

$$(3.28) \quad \begin{aligned} b[\boldsymbol{\eta}_h, p_h] & = -[[\partial_t \boldsymbol{\xi}_h, \boldsymbol{\eta}_h]] - a_1[\mathbf{v}_h, \boldsymbol{\eta}_h] \\ & - a_2[\mathbf{u}_h, \boldsymbol{\eta}_h] + [[\mathbf{f}, \boldsymbol{\eta}_h]] \quad \forall \boldsymbol{\eta}_h \in X^h, \forall t \in [0, T]. \end{aligned}$$

To show the existence of a $p_h \in C([0, T]; Q_1^h)$ satisfying (3.27) or (3.28), we need to verify a discrete inf-sup condition for $b[\cdot, \cdot]$, which will be presented below; this will be the task of this subsection. To derive an estimate for p_h , we need an estimate for $\|\partial_t \boldsymbol{\xi}_h\|_{0,\Omega}$, or $\|\partial_t \mathbf{v}_h\|_{0,\Omega_1}$ and $\|\partial_{tt} \mathbf{u}_h\|_{0,\Omega_2}$; these will be derived in section 3.4.

The inf-sup condition we will verify is

$$(3.29) \quad \inf_{q_h \in Q_1^h} \sup_{\boldsymbol{\eta}_h \in X^h} \frac{b[\boldsymbol{\eta}_h, q_h]}{\|\boldsymbol{\eta}_h\|_{1,\Omega} \|q_h\|_{0,\Omega_1}} \geq C.$$

This inf-sup condition was proved in [2] for a special choice of X_h and Q_1^h . We will establish (3.29) for the general case under assumption (3.12). To this end, we will

first need the following lemma, and we will need to prove the inf-sup condition

$$(3.30) \quad \inf_{q_h \in Q_1^h} \sup_{\mathbf{v}_h \in X_1^h} \frac{b[\mathbf{v}_h, q_h]}{\|\mathbf{v}_h\|_{1, \Omega_1} \|q_h\|_{0, \Omega_1}} \geq C.$$

LEMMA 3.3. *For each constant d , there exists a piecewise linear function $\mathbf{v} \in X_1^{h_0}$ such that*

$$\int_{\Gamma_0} \mathbf{v} \cdot \mathbf{n} \, d\Gamma = -d, \quad b[\mathbf{v}, d] = |d|^2, \quad \text{and} \quad \|\mathbf{v}\|_{1, \Omega_1} \leq C|d|,$$

where \mathbf{n} denotes the unit outward-pointing normal along $\partial\Omega_1$, and the constant C depends only on the coarse triangulation $\mathcal{T}^{h_0}(\Omega_1)$.

Proof. We give the complete proof for the two-dimensional case and discuss the ideas for the three-dimensional case in an ensuing remark.

We choose from $\mathcal{T}^{h_0}(\Omega)$ a layer of triangles $K \equiv \cup_{j=1}^{J_0} K_j \subset \bar{\Omega}_1$ adjacent to Γ_0 , i.e., each K_j has either a side or a vertex on Γ_0 . We denote the vertices on $\Gamma_0 \cap \partial K$ by $A_j, j = 0, 1, \dots, J_0$. We define the C^0 , piecewise linear vector function $\mathbf{v} = (v_1, v_2)$ on K as follows:

$$\left\{ \begin{array}{l} \mathbf{v} = \mathbf{0} \text{ at points } A_0 \text{ and } A_{J_0}, \\ \mathbf{v} = \mathbf{0} \text{ at all vertices of } K \text{ belonging to the interior of } \Omega_1, \\ \mathbf{v} \cdot \mathbf{n}_{j-1} = -\bar{d} \text{ and } \mathbf{v} \cdot \mathbf{n}_j = -\bar{d} \text{ at } A_j, \quad j = 1, \dots, J_0 - 1, \mathbf{n}_{j-1} \neq \mathbf{n}_j, \\ \mathbf{v} \cdot \mathbf{n}_{j-1} = -\bar{d} \text{ and } \mathbf{v} \cdot \boldsymbol{\tau}_j = 0 \text{ at } A_j, \quad j = 1, \dots, J_0 - 1, \mathbf{n}_{j-1} = \mathbf{n}_j, \end{array} \right.$$

where

$$\bar{d} = d \left/ \left(\frac{|A_0 A_1|}{2} + \sum_{j=2}^{J_0-1} |A_{j-1} A_j| + \frac{|A_{J_0-1} A_{J_0}|}{2} \right) \right.$$

and \mathbf{n}_j and $\boldsymbol{\tau}_j$ denote the unit, outward-pointing normal and unit tangent vectors, respectively, on $\partial\Omega_1 \cap \bar{A}_{j-1} A_j$. Note that \mathbf{n}_j and $\boldsymbol{\tau}_j$ are defined with respect to the segment $\bar{A}_{j-1} A_j$ so that they are well defined. Clearly, the values of $v_1(A_j)$ and $v_2(A_j)$ are proportional to \bar{d} . We can write

$$v_i(\mathbf{x}) = \sum_{j=1}^{J_0-1} v_i(A_j) L_j^{h_0}(\mathbf{x}), \quad i = 1, 2,$$

where for each j , $L_j^{h_0}(\mathbf{x})$ is the continuous piecewise linear basis function (the shape function) associated with the vertex A_j . Then,

$$\|v_i\|_{1, K}^2 \leq C \sum_{j=1}^{J_0-1} |v_i(A_j)|^2 \|L_j^{h_0}\|_{1, K}^2 \leq C|\bar{d}|^2 \sum_{j=1}^{J_0-1} \|L_j^{h_0}\|_{1, K}^2$$

so that

$$\|\mathbf{v}\|_{1, K} \leq C|\bar{d}|.$$

We extend \mathbf{v} to Ω_1 by zero outside K and denote the extended function still by \mathbf{v} . Then we readily have $\mathbf{v} \in X_1^{h_0}$,

$$\|\mathbf{v}\|_{1,\Omega_1} = \|\mathbf{v}\|_{1,K} \leq C|\bar{d}| \leq C|d|,$$

and

$$\begin{aligned} \int_{\Gamma_0} \mathbf{v} \cdot \mathbf{n} d\Gamma &= \sum_{j=1}^{J_0} \int_{A_{j-1}A_j} \mathbf{v} \cdot \mathbf{n} d\Gamma \\ &= -\bar{d} \left(\frac{|A_0A_1|}{2} + \sum_{j=2}^{J_0-1} |A_{j-1}A_j| + \frac{|A_{J_0-1}A_{J_0}|}{2} \right) = -d. \end{aligned}$$

Using Green's theorem and the last equality, we have

$$b[\mathbf{v}, d] = -d \int_{\Omega_1} \nabla \cdot \mathbf{v} d\Omega = -d \int_{\Gamma_0} \mathbf{v} \cdot \mathbf{n} d\Gamma = d^2. \quad \square$$

Remark 1. In the three-dimensional case we merely need assume that $[\mathcal{T}^{h_0}(\Omega)]|_{\Gamma_0}$ contains a vertex P_0 shared by exactly three triangles. Indeed, in forming the coarse triangulation $\mathcal{T}^{h_0}(\Omega)$, we may simply choose a partition on a flat piece of Γ_0 to meet this requirement. Then, we define a \mathbf{v} to satisfy $\mathbf{v} \cdot \mathbf{n} = \bar{d}$ and $\mathbf{v} \times \mathbf{n} = \mathbf{0}$ at P_0 , and $\mathbf{v} = \mathbf{0}$ at all other vertices, where \bar{d} is a suitable scaling of d .

Next we prove inf-sup condition (3.30) based on the inf-sup assumption (3.12) for the pair $\{\tilde{X}_1^h, M_1^h\} \equiv \{X_1^h \cap \mathbf{H}_0^1(\Omega_1), Q_1^h \cap L_0^2(\Omega_1)\}$.

THEOREM 3.4. *The pair $\{X_1^h, Q_1^h\}$ satisfies inf-sup condition (3.30).*

Proof. Owing to [19, Remark 1.4, p. 118], the inf-sup condition (3.30) is equivalent to

$$(3.31) \quad \begin{aligned} \forall q_h \in Q_1^h, \quad \text{there exists } \mathbf{v}_h \in X_1^h \quad \text{such that} \\ b[\mathbf{v}_h, q_h] \geq C\|q_h\|_{0,\Omega_1}^2 \quad \text{and} \quad \|\mathbf{v}_h\|_{1,\Omega_1} \leq C\|q_h\|_{0,\Omega_1}. \end{aligned}$$

Let $q_h \in Q_1^h$ be given. Set

$$\bar{q}_h = \frac{1}{|\Omega_1|} \int_{\Omega_1} q_h d\Omega \quad \text{and} \quad \tilde{q}_h = q_h - \bar{q}_h.$$

Then $q_h = \tilde{q}_h + \bar{q}_h$ in Ω_1 and $\|q_h\|_{0,\Omega_1}^2 = \|\tilde{q}_h\|_{0,\Omega_1}^2 + \|\bar{q}_h\|_{0,\Omega_1}^2$. Obviously, $\tilde{q}_h \in M_1^h \equiv Q_1^h \cap L_0^2(\Omega_1)$ so that, by inf-sup condition (3.12) for the pair $\{\tilde{X}_1^h, M_1^h\}$, we may choose a $\tilde{\mathbf{v}}_h \in \tilde{X}_1^h$ such that

$$b[\tilde{\mathbf{v}}_h, \tilde{q}_h] = \|\tilde{q}_h\|_{0,\Omega_1}^2 \quad \text{and} \quad \|\tilde{\mathbf{v}}_h\|_{1,\Omega_1} \leq C\|\tilde{q}_h\|_{0,\Omega_1}.$$

By Lemma 3.3 with $d = \|\bar{q}_h\|_{0,\Omega_1}$, we may choose a $\bar{\mathbf{v}}_h \in X_1^h$ such that

$$b[\bar{\mathbf{v}}_h, \bar{q}_h] = \|\bar{q}_h\|_{0,\Omega_1}^2 \quad \text{and} \quad \|\bar{\mathbf{v}}_h\|_{1,\Omega_1} \leq C\|\bar{q}_h\|_{0,\Omega_1}.$$

(We recall that we assumed that $\mathcal{T}^h(\Omega_1)$ is a refinement of a coarse triangulation $\mathcal{T}^{h_0}(\Omega_1)$ so that a piecewise linear function on $\mathcal{T}^{h_0}(\Omega_1)$ belongs to X_1^h .) Setting

$\mathbf{v}_h = \tilde{\mathbf{v}}_h + \alpha \bar{\mathbf{v}}_h$ for some $\alpha > 0$ (to be determined), we have

$$\begin{aligned} b[\mathbf{v}_h, q_h] &= b[\tilde{\mathbf{v}}_h, \tilde{q}_h] + b[\tilde{\mathbf{v}}_h, \bar{q}_h] + \alpha b[\bar{\mathbf{v}}_h, \tilde{q}_h] + \alpha b[\bar{\mathbf{v}}_h, \bar{q}_h] \\ &\geq \|\tilde{q}_h\|_{0,\Omega_1}^2 + 0 - C\alpha \|\tilde{q}_h\|_{0,\Omega_1} \|\bar{\mathbf{v}}_h\|_{1,\Omega_1} + \alpha \|\bar{q}_h\|_{0,\Omega_1}^2 \\ &\geq \|\tilde{q}_h\|_{0,\Omega_1}^2 - C\alpha \|\tilde{q}_h\|_{0,\Omega_1} \|\bar{q}_h\|_{0,\Omega_1} + \alpha \|\bar{q}_h\|_{0,\Omega_1}^2 \\ &\geq \|\tilde{q}_h\|_{0,\Omega_1}^2 - [C\alpha \|\tilde{q}_h\|_{0,\Omega_1}^2 + \frac{\alpha}{2} \|\bar{q}_h\|_{0,\Omega_1}^2] + \alpha \|\bar{q}_h\|_{0,\Omega_1}^2 \\ &= (1 - C\alpha) \|\tilde{q}_h\|_{0,\Omega_1}^2 + \frac{\alpha}{2} \|\bar{q}_h\|_{0,\Omega_1}^2 \end{aligned}$$

so that by choosing a sufficiently small $\alpha > 0$ we obtain

$$b[\mathbf{v}_h, q_h] \geq \min\{1 - C\alpha, \alpha/2\} \left(\|\tilde{q}_h\|_{0,\Omega_1}^2 + \frac{1}{2} \|\bar{q}_h\|_{0,\Omega_1}^2 \right) \geq C \|q_h\|_{0,\Omega_1}^2.$$

Also,

$$\|\mathbf{v}_h\|_{1,\Omega_1} \leq \|\tilde{\mathbf{v}}_h\|_{1,\Omega_1} + \|\bar{\mathbf{v}}_h\|_{1,\Omega_1} \leq C \|\tilde{q}_h\|_{0,\Omega_1} + C \|\bar{q}_h\|_{0,\Omega_1} \leq C \|q_h\|_{0,\Omega_1}.$$

Hence, we have proved (3.31) which is equivalent to (3.30). \square

We now prove inf-sup condition (3.29) for $\{X^h, Q_1^h\}$.

THEOREM 3.5. $\{X^h, Q_1^h\}$ satisfies the inf-sup condition (3.29).

Proof. Let the discrete extension operator $E^h : X_1^h \rightarrow X^h$ be defined as follows: for any $\mathbf{v}_h \in X_1^h$, $(E^h \mathbf{v}_h)|_{\bar{\Omega}_1} = \mathbf{v}_h$ and $(E^h \mathbf{v}_h)|_{\Omega_2} \in X_2^h$ is the solution of

$$[\nabla(E^h \mathbf{v}_h), \nabla \mathbf{z}_h]_{\Omega_2} = 0 \quad \forall \mathbf{z}_h \in X_2^h \cap \mathbf{H}_0^1(\Omega_2), \quad (E^h \mathbf{v}_h)|_{\Gamma_2} = \mathbf{0}, \quad (E^h \mathbf{v}_h)|_{\Gamma_0} = \mathbf{v}_h|_{\Gamma_0}.$$

It is well known (see, e.g., [23] and [1]) that $\|E^h \mathbf{v}_h\|_{1,\Omega_2} \leq C \|\mathbf{v}_h\|_{1/2,\Gamma_0}$ so that

$$\begin{aligned} \|E^h \mathbf{v}_h\|_{1,\Omega} &\leq C (\|(E^h \mathbf{v}_h)|_{\Omega_1}\|_{1,\Omega_1} + \|(E^h \mathbf{v}_h)|_{\Omega_2}\|_{1,\Omega_2}) \\ &\leq C (\|\mathbf{v}_h\|_{1,\Omega_1} + \|\mathbf{v}_h\|_{1/2,\Gamma_0}) \leq C \|\mathbf{v}_h\|_{1,\Omega_1} \quad \forall \mathbf{v}_h \in X_1^h. \end{aligned}$$

Then, for every $q_h \in Q_1^h$ we have

$$\begin{aligned} \sup_{\boldsymbol{\eta}_h \in X^h} \frac{b[\boldsymbol{\eta}_h, q_h]}{\|q_h\|_{0,\Omega_1} \|\boldsymbol{\eta}_h\|_{1,\Omega}} &\geq \sup_{\mathbf{v}_h \in X_1^h} \frac{b[E^h \mathbf{v}_h, q_h]}{\|q_h\|_{0,\Omega_1} \|E^h \mathbf{v}_h\|_{1,\Omega}} \\ &\geq C \sup_{\mathbf{v}_h \in X_1^h} \frac{b[E^h \mathbf{v}_h, q_h]}{\|q_h\|_{0,\Omega_1} \|\mathbf{v}_h\|_{1,\Omega_1}} = C \sup_{\mathbf{v}_h \in X_1^h} \frac{b[\mathbf{v}_h, q_h]}{\|q_h\|_{0,\Omega_1} \|\mathbf{v}_h\|_{1,\Omega_1}} \geq C, \end{aligned}$$

where the last step is valid because of (3.30). \square

As a direct consequence of [19, Lemma 4.1, p. 58], Theorem 3.8, and the inf-sup condition (3.29), we obtain the following theorem concerning the existence of a discrete pressure. Note that an estimate for p_h will be established in section 3.4 only after we have derived strong energy estimates, particularly the estimate for $\|\partial_t \boldsymbol{\xi}_h\|_{L^2(0,T;L^2(\Omega))}$.

THEOREM 3.6. Assume that $\mathbf{f}_1, \mathbf{v}_0, \mathbf{f}_2, \mathbf{u}_0$, and \mathbf{u}_1 satisfy (2.6), and let $\boldsymbol{\xi}_h \in C^1([0, T]; \boldsymbol{\Psi}^h)$ be the solution of (3.21)–(3.22). Let $(\mathbf{v}_h, \mathbf{u}_h) \in C^1([0, T]; X^h|_{\Omega_1}) \times C^1([0, T]; X_2^h)$ be the solution of (3.25) and (3.15)–(3.19). Then there exists a unique $p_h \in C([0, T]; Q_1^h)$ satisfying (3.28) and (3.15).

Proof. The existence and uniqueness of a $p_h \in C([0, T]; Q_1^h)$ satisfying (3.28) follow directly from [19, Lemma 4.1, p. 58], Theorem 3.8, and the inf-sup condition (3.29). Since (3.28) is equivalent to (3.27), we also conclude that p_h satisfies (3.27) and is the unique such solution. \square

3.4. Strong a priori energy estimates for the finite element solutions.

In the finite element system (3.25) and (3.15)–(3.19) the discrete initial conditions are arbitrary approximations of the corresponding continuous initial data. We now make a particular choice of discrete initial data that will allow us to derive an estimate for $\|\partial_t \boldsymbol{\xi}_h\|_{0,\Omega}$ under additional assumptions on the data. Such an estimate can then be used to derive an estimate for $\|p_h\|_{L^2(0,T;L^2(\Omega_1))}$. (The existence of a discrete pressure p_h satisfying (3.14) was shown in section 3.3.) The estimates on p_h and $\partial_t \boldsymbol{\xi}_h$ will be needed in order to prove the convergence of finite element solutions, since finite element formulations involve the term $b[\boldsymbol{\eta}_h, p_h]$, which, in general, does not vanish for $\boldsymbol{\eta}_h \in X^h$.

We first study the approximation of the initial condition. We choose $(\mathbf{v}_{0,h}, \mathbf{u}_{1,h}) \in \boldsymbol{\Psi}^h$ and $p_{0,h} \in Q_1^h$ to be the solution of

$$(3.32) \quad \begin{aligned} & a_1[\mathbf{v}_{0,h}, \boldsymbol{\eta}_h] + [\mathbf{u}_{1,h}, \boldsymbol{\eta}_h]_{\Omega_2} + b[\boldsymbol{\eta}_h, p_{0,h}] \\ & = a_1[\mathbf{v}_0, \boldsymbol{\eta}_h] + [\mathbf{u}_1, \boldsymbol{\eta}_h]_{\Omega_2} + b[\boldsymbol{\eta}_h, p_0] \quad \forall \boldsymbol{\eta}_h \in X^h, \end{aligned}$$

$$(3.33) \quad b[\mathbf{v}_{0,h}, q_h] = 0 \quad \forall q_h \in Q_1^h \quad \text{and} \quad \mathbf{v}_{0,h}|_{\Gamma_0} = \mathbf{u}_{1,h}|_{\Gamma_0},$$

where p_0 is the initial pressure field associated with the initial velocity field \mathbf{v}_0 .

LEMMA 3.7. *Assume that $\mathbf{v}_0 \in X_1$, $p_0 \in L^2(\Omega_1)$, $\mathbf{u}_1 \in X_2$, and $\mathbf{v}_0|_{\Gamma_0} = \mathbf{u}_1|_{\Gamma_0}$. Then there exists a unique triplet $(\mathbf{v}_{0,h}, p_{0,h}, \mathbf{u}_{1,h}) \in X_1^h \times Q_1^h \times X_2^h$ which satisfies (3.32)–(3.33) and*

$$(3.34) \quad \begin{aligned} & \|\mathbf{v}_{0,h} - \mathbf{v}_0\|_{1,\Omega_1} + \|\mathbf{u}_{1,h} - \mathbf{u}_1\|_{0,\Omega_2} + \|p_{0,h} - p_0\|_{0,\Omega_1} \\ & \leq C(\|\boldsymbol{\eta}_h - \mathbf{v}_0\|_{1,\Omega_1} + \|\boldsymbol{\eta}_h - \mathbf{u}_1\|_{0,\Omega_2} + \|q_h - p_0\|_{0,\Omega_1}) \quad \forall (\boldsymbol{\eta}_h, q_h) \in X^h \times Q_1^h. \end{aligned}$$

If, in addition, $\mathbf{v}_0 \in \mathbf{H}^{r+1}(\Omega_1)$, $p_0 \in H^r(\Omega_1)$, and $\mathbf{u}_1 \in \mathbf{H}^{r+1}(\Omega_2)$ for some $r \in [0, k]$ (k being the integer appearing in the approximation properties), then

$$(3.35) \quad \begin{aligned} & \|\mathbf{v}_{0,h} - \mathbf{v}_0\|_{1,\Omega_1} + \|\mathbf{u}_{1,h} - \mathbf{u}_1\|_{0,\Omega_2} + \|p_{0,h} - p_0\|_{0,\Omega_1} \\ & \leq Ch^r(\|\mathbf{v}_0\|_{r+1,\Omega_1} + \|\mathbf{u}_1\|_{r+1,\Omega_2} + \|p_0\|_{r,\Omega_1}). \end{aligned}$$

Proof. We set $\tilde{X} = \{\boldsymbol{\eta} \in \mathbf{L}^2(\Omega) : \boldsymbol{\eta}|_{\Omega_1} \in X_1, \operatorname{div} \boldsymbol{\eta}|_{\Omega_1} = 0\}$ and equip \tilde{X} with the inner product

$$[\boldsymbol{\xi}, \boldsymbol{\eta}]_{\tilde{X}} = a_1[\boldsymbol{\xi}, \boldsymbol{\eta}] + [\boldsymbol{\xi}, \boldsymbol{\eta}]_{\Omega_2} \quad \forall \boldsymbol{\xi}, \boldsymbol{\eta} \in \tilde{X}.$$

It is easy to check that \tilde{X} is a Hilbert space. The continuous inf-sup condition (2.4) implies

$$\begin{aligned} \inf_{q \in L^2(\Omega_1)} \sup_{\boldsymbol{\eta} \in \tilde{X}} \frac{b[\boldsymbol{\eta}, q]}{\|\boldsymbol{\eta}\|_{\tilde{X}} \|q\|_{0,\Omega_1}} & \geq \inf_{q \in L^2(\Omega_1)} \sup_{\boldsymbol{\eta} \in \mathbf{H}_0^1(\Omega)} \frac{b[\boldsymbol{\eta}, q]}{\|\boldsymbol{\eta}\|_{\tilde{X}} \|q\|_{0,\Omega_1}} \\ & \geq \inf_{q \in L^2(\Omega_1)} \sup_{\boldsymbol{\eta} \in \mathbf{H}_0^1(\Omega)} \frac{b[\boldsymbol{\eta}, q]}{\|\boldsymbol{\eta}\|_{1,\Omega} \|q\|_{0,\Omega_1}} \geq C. \end{aligned}$$

Thus, by [19, Theorem 1.1, p. 114], there exists a unique $(\tilde{\boldsymbol{\xi}}_0, \tilde{p}_0) \in \tilde{X} \times L^2(\Omega_1)$ satisfying

$$(3.36) \quad [\tilde{\boldsymbol{\xi}}_0, \boldsymbol{\eta}]_{\tilde{X}} + b[\boldsymbol{\eta}, \tilde{p}_0] = a_1[\mathbf{v}_0, \boldsymbol{\eta}] + [\mathbf{u}_1, \boldsymbol{\eta}]_{\Omega_2} + b[\boldsymbol{\eta}, p_0] \quad \forall \boldsymbol{\eta} \in \tilde{X},$$

$$(3.37) \quad b[\tilde{\boldsymbol{\xi}}_0, q] = 0 \quad \forall q \in L^2(\Omega_1).$$

As $\boldsymbol{\xi}_0$ defined by (2.11) and p_0 constitute an obvious solution to (3.36)–(3.37), we have

$$(3.38) \quad \tilde{\boldsymbol{\xi}}_0 = \boldsymbol{\xi}_0 = \begin{cases} \boldsymbol{\xi}|_{\Omega_1} = \mathbf{v}_0, \\ \boldsymbol{\xi}|_{\Omega_2} = \mathbf{u}_1, \end{cases} \quad \text{and} \quad \tilde{p}_0 = p_0.$$

Similarly, the discrete inf-sup condition (3.29) implies

$$\inf_{q_h \in Q_1^h} \sup_{\boldsymbol{\eta}_h \in X^h} \frac{b[\boldsymbol{\eta}_h, q_h]}{\|\boldsymbol{\eta}_h\|_{\bar{X}} \|q_h\|_{0, \Omega_1}} \geq \inf_{q_h \in Q_1^h} \sup_{\boldsymbol{\eta}_h \in X^h} \frac{b[\boldsymbol{\eta}_h, q_h]}{\|\boldsymbol{\eta}_h\|_{1, \Omega} \|q_h\|_{0, \Omega_1}} \geq C,$$

so that by [19, Theorem 1.1, p. 114] there exists a unique $(\boldsymbol{\xi}_{0,h}, p_{0,h}) \in X^h \times Q_1^h$ satisfying

$$(3.39) \quad [\boldsymbol{\xi}_{0,h}, \boldsymbol{\eta}_h]_{\bar{X}} + b[\boldsymbol{\eta}_h, p_{0,h}] = a_1[\mathbf{v}_0, \boldsymbol{\eta}_h] + [\mathbf{u}_1, \boldsymbol{\eta}_h]_{\Omega_2} + b[\boldsymbol{\eta}_h, p_0] \quad \forall \boldsymbol{\eta}_h \in X^h,$$

$$(3.40) \quad b[\boldsymbol{\xi}_{0,h}, q_h] = 0 \quad \forall q_h \in Q_1^h;$$

moreover, the following error estimate holds:

$$(3.41) \quad \|\boldsymbol{\xi}_{0,h} - \boldsymbol{\xi}_0\|_{\bar{X}} + \|p_{0,h} - p_0\|_{0, \Omega_1} \leq C(\|\boldsymbol{\eta}_h - \boldsymbol{\xi}_0\|_{\bar{X}} + \|q_h - p_0\|_{0, \Omega_1}) \quad \forall (\boldsymbol{\eta}_h, q_h) \in X^h \times Q_1^h.$$

By setting

$$(3.42) \quad \mathbf{v}_{0,h} = \boldsymbol{\xi}_{0,h}|_{\Omega_1} \quad \text{and} \quad \mathbf{u}_{1,h} = \boldsymbol{\xi}_{0,h}|_{\Omega_2},$$

we see that (3.41) is equivalent to (3.34) and that (3.32)–(3.33) are satisfied. The uniqueness of the solution $(\mathbf{v}_{0,h}, p_{0,h}, \mathbf{u}_{1,h})$ for (3.32)–(3.33) follows from the uniqueness of the solution $(\boldsymbol{\xi}_{0,h}, p_{0,h})$ for (3.39)–(3.40).

Next, assuming that $\mathbf{v}_0 \in \mathbf{H}^{r+1}(\Omega_1)$, $p_0 \in H^r(\Omega_1)$, and $\mathbf{u}_1 \in \mathbf{H}^{r+1}(\Omega_2)$ for some $r \in [1, k]$, we proceed to prove (3.35) by making a particular choice of $\boldsymbol{\eta}_h$ in (3.34). Let $(\bar{\mathbf{v}}_{0,h}, \bar{p}_{0,h}) \in X_1^h \times Q_1^h$ be the unique finite element solution of the following Stokes system on Ω_1 :

$$a_1[\bar{\mathbf{v}}_{0,h}, \mathbf{z}_h] + b[\mathbf{z}_h, \bar{p}_{0,h}] = a_1[\mathbf{v}_0, \mathbf{z}_h] + b[\mathbf{z}_h, \bar{p}_0] \quad \forall \mathbf{z}_h \in X_1^h \cap \mathbf{H}_0^1(\Omega_1),$$

$$b[\bar{\mathbf{v}}_{0,h}, q_h] = 0 \quad \forall q_h \in Q_1^h \cap L_0^2(\Omega_1),$$

$$\bar{\mathbf{v}}_{0,h}|_{\Gamma_1} = \mathbf{0} \quad \text{and} \quad [\bar{\mathbf{v}}_{0,h}, \mathbf{s}_h]_{0, \Gamma_0} = [\mathbf{v}_0, \mathbf{s}_h]_{0, \Gamma_0} \quad \forall \mathbf{s}_h \in X_1^h|_{\Gamma_0},$$

where $\bar{p}_0 = p_0 - (1/|\Omega_1|) \int_{\Omega_1} p_0 \, dx$. Using the results of [23] concerning error estimates for the finite element approximations of the Stokes equations with inhomogeneous boundary conditions, we obtain

$$(3.43) \quad \|\bar{\mathbf{v}}_{0,h} - \mathbf{v}_0\|_{1, \Omega_1} + \|\bar{p}_{0,h} - \bar{p}_0\|_{0, \Omega_1} \leq Ch^r (\|\mathbf{v}_0\|_{r+1, \Omega_1} + \|\bar{p}_0\|_{r, \Omega_1}) \leq Ch^r (\|\mathbf{v}_0\|_{r+1, \Omega_1} + \|p_0\|_{r, \Omega_1}).$$

Analogously, let $\bar{\mathbf{u}}_{1,h} \in X_2^h$ be the unique finite element solution of the following elliptic system on Ω_2 with an inhomogeneous boundary condition:

$$(3.44) \quad \begin{aligned} [\nabla \bar{\mathbf{u}}_{1,h}, \nabla \mathbf{w}_h]_{\Omega_2} &= [\nabla \mathbf{u}_1, \nabla \mathbf{w}_h]_{\Omega_2} \quad \forall \mathbf{w}_h \in X_2^h \cap \mathbf{H}_0^1(\Omega_2), \\ \bar{\mathbf{u}}_{1,h}|_{\Gamma_2} &= \mathbf{0} \quad \text{and} \quad [\bar{\mathbf{u}}_{1,h}, \mathbf{s}_h]_{0,\Gamma_0} = [\mathbf{u}_1, \mathbf{s}_h]_{0,\Gamma_0} \quad \forall \mathbf{s}_h \in X_2^h|_{\Gamma_0}. \end{aligned}$$

Then we have

$$(3.45) \quad \|\bar{\mathbf{u}}_{1,h} - \mathbf{u}_1\|_{0,\Omega_2} \leq \|\bar{\mathbf{u}}_{1,h} - \mathbf{u}_1\|_{1,\Omega_2} \leq Ch^r \|\mathbf{u}_1\|_{r+1,\Omega_2}.$$

The assumption $\mathbf{v}_0|_{\Gamma_0} = \mathbf{u}_1|_{\Gamma_0}$ implies $\bar{\mathbf{v}}_{0,h}|_{\Gamma_0} = \bar{\mathbf{u}}_{1,h}|_{\Gamma_0}$, so that the element $\bar{\boldsymbol{\eta}}_h$ defined by

$$\bar{\boldsymbol{\eta}}_h|_{\Omega_1} = \begin{cases} \bar{\mathbf{v}}_{0,h} & \text{in } \Omega_1, \\ \bar{\mathbf{u}}_{1,h} & \text{in } \Omega_2 \end{cases}$$

satisfies $\bar{\boldsymbol{\eta}}_h \in X^h$. By choosing $\boldsymbol{\eta}_h = \bar{\boldsymbol{\eta}}_h$ and $q_h = \bar{p}_{0,h} + (1/|\Omega_1|) \int_{\Omega_1} p_0 \, d\mathbf{x}$ in (3.34) and using (3.43)–(3.45), we arrive at (3.35). \square

We now derive a strong a priori energy estimate for the auxiliary finite element solution $\boldsymbol{\xi}_h$.

THEOREM 3.8. *Assume that $\mathbf{f}_1, \mathbf{v}_0, \mathbf{f}_2, \mathbf{u}_0$, and \mathbf{u}_1 satisfy (2.6) and*

$$(3.46) \quad \partial_t \mathbf{f}_i \in L^2(0, T; \mathbf{L}^2(\Omega_i)), \quad i = 1, 2, \quad \mathbf{v}_0 \in \mathbf{H}^2(\Omega_1), \quad \mathbf{u}_1 \in \mathbf{H}^2(\Omega_1), \quad \mathbf{u}_0 \in \mathbf{H}^2(\Omega_2).$$

Assume further that there exists a $p_0 \in H^1(\Omega_1)$ such that

$$(3.47) \quad (p_0 \mathbf{n}_1 - \mu_1 \nabla \mathbf{v}_0 \cdot \mathbf{n}_1)|_{\Gamma_0} = (\mu_2 \nabla \mathbf{u}_0 \cdot \mathbf{n}_2 + (\lambda_2 + \mu_2)(\operatorname{div} \mathbf{u}_0) \mathbf{n}_2)|_{\Gamma_0},$$

where \mathbf{n}_i denotes the outward-pointing normal along $\partial\Omega_i$, $i = 1, 2$. Then there exists a unique solution $\boldsymbol{\xi}_h \in C^1([0, T]; \boldsymbol{\Psi}^h)$ for (3.21)–(3.22) with the initial condition $\boldsymbol{\xi}_{0,h}$ defined by

$$(3.48) \quad \boldsymbol{\xi}_{0,h}|_{\Omega_1} = \mathbf{v}_{0,h} \quad \text{and} \quad \boldsymbol{\xi}_{0,h}|_{\Omega_2} = \mathbf{u}_{1,h},$$

where $\mathbf{v}_{0,h}$ and $\mathbf{u}_{1,h}$ are determined by (3.32)–(3.33). Moreover, $\boldsymbol{\xi}_h$ satisfies the estimates

$$(3.49) \quad \begin{aligned} &\|\boldsymbol{\xi}_h\|_{L^2(0,T;\mathbf{L}^2(\Omega))}^2 + \|\boldsymbol{\xi}_h\|_{L^2(0,T;\mathbf{H}^1(\Omega_1))}^2 + \left\| \int_0^t \boldsymbol{\xi}_h(s) \, ds \right\|_{L^\infty(0,T;\mathbf{H}^1(\Omega_2))}^2 \\ &\leq Ce^{CT} (\|\mathbf{f}\|_{L^2(0,T;\mathbf{L}^2(\Omega))}^2 + \|\mathbf{u}_0\|_{1,\Omega_2}^2 + \|\mathbf{v}_0\|_{1,\Omega_1}^2 + \|\mathbf{u}_1\|_{0,\Omega_2}^2 + \|p_0\|_{0,\Omega_1}^2) \end{aligned}$$

and

$$(3.50) \quad \begin{aligned} &\|\partial_t \boldsymbol{\xi}_h\|_{L^\infty(0,T;\mathbf{L}^2(\Omega))}^2 + \|\partial_t \boldsymbol{\xi}_h\|_{L^2(0,T;\mathbf{H}^1(\Omega_1))}^2 + \left\| \int_0^t \partial_t \boldsymbol{\xi}_h(s) \, ds \right\|_{L^\infty(0,T;\mathbf{H}^1(\Omega_2))}^2 \\ &\leq Ce^{CT} (\|\mathbf{f}\|_{H^1(0,T;\mathbf{L}^2(\Omega))}^2 + \|\mathbf{u}_0\|_{2,\Omega_2}^2 + \|\mathbf{v}_0\|_{2,\Omega_1}^2 + \|p_0\|_{1,\Omega_1}^2 + \|\mathbf{u}_1\|_{2,\Omega_2}^2). \end{aligned}$$

Proof. By Theorem 3.1, there exists a unique solution $\boldsymbol{\xi}_h \in C^1([0, T]; \boldsymbol{\Psi}^h)$ for (3.21)–(3.22) and (3.24). We note that, by virtue of Lemma 3.7, the initial condition $\boldsymbol{\xi}_{0,h} \in \boldsymbol{\Psi}^h$ satisfies the estimate

$$\|\boldsymbol{\xi}_{0,h}\|_{1,\Omega_1} + \|\boldsymbol{\xi}_{0,h}\|_{0,\Omega_2} \leq C(\|\mathbf{v}_0\|_{1,\Omega_1} + \|p_0\|_{0,\Omega_1} + \|\mathbf{u}_1\|_{0,\Omega_2}).$$

Thus (3.49) follows from the last estimate and (3.24).

Defining $\zeta_h = \partial_t \xi_h$ and differentiating (3.21), we obtain that for each $t \in [0, T]$

$$(3.51) \quad \begin{aligned} & \rho_1[\partial_t \zeta_h, \boldsymbol{\eta}_h]_{\Omega_1} + \rho_2[\partial_t \zeta_h, \boldsymbol{\eta}_h]_{\Omega_2} + a_1[\zeta_h, \boldsymbol{\eta}_h] + a_2 \left[\int_0^t \zeta_h(s) ds, \boldsymbol{\eta}_h \right] \\ & = \rho_1[\partial_t \mathbf{f}_1, \boldsymbol{\eta}_h]_{\Omega_1} + \rho_2[\partial_t \mathbf{f}_2, \boldsymbol{\eta}_h]_{\Omega_2} - a_2[\xi_h(0), \boldsymbol{\eta}_h] \quad \forall \boldsymbol{\eta}_h \in \boldsymbol{\Psi}^h. \end{aligned}$$

Setting $\boldsymbol{\eta}_h = \zeta_h(t)$ in (3.51) and integrating in t , we obtain

$$\begin{aligned} & [[\zeta_h(t), \zeta_h(t)]] + \int_0^t a_1[\zeta_h(s), \zeta_h(s)] ds + a_2 \left[\int_0^t \zeta_h(s) ds, \int_0^t \zeta_h(s) ds \right] \\ & \leq C(\|\zeta_h(0)\|_{0,\Omega}^2 + \|\mathbf{f}\|_{L^2(0,T;\mathbf{L}^2(\Omega))}^2) + a_2 \left[\mathbf{u}_0, \int_0^t \zeta_h(s) ds \right] + \int_0^t \|\zeta_h(s)\|_{0,\Omega}^2 ds \\ & \leq C(\|\zeta_h(0)\|_{0,\Omega}^2 + \|\mathbf{f}\|_{L^2(0,T;\mathbf{L}^2(\Omega))}^2 + \|\mathbf{u}_0\|_{1,\Omega_2}^2) \\ & \quad + \frac{1}{2} a_2 \left[\int_0^t \zeta_h(s) ds, \int_0^t \zeta_h(s) ds \right] + \int_0^t \|\zeta_h(s)\|_{0,\Omega}^2 ds, \end{aligned}$$

so that

$$(3.52) \quad \begin{aligned} & \|\zeta_h(t)\|_{0,\Omega}^2 + \int_0^t a_1[\zeta_h(t), \zeta_h(t)] dt + a_2 \left[\int_0^t \zeta_h(s) ds, \int_0^t \zeta_h(s) ds \right] \\ & \leq C(\|\zeta_h(0)\|_{0,\Omega}^2 + \|\partial_t \mathbf{f}\|_{L^2(0,T;\mathbf{L}^2(\Omega))}^2 + \|\xi_h(0)\|_{1,\Omega_2}^2) + \int_0^t \|\zeta_h(s)\|_{0,\Omega}^2 ds. \end{aligned}$$

Dropping the second and third terms on the left-hand side of (3.52) and then applying the following version of Gronwall's inequality [12, p. 625],

$$(3.53) \quad \text{if } r(t) \leq C_1 + C_2 \int_0^t r(s) ds, \text{ then } r(t) \leq C_1(1 + C_2 t)e^{C_2 t},$$

we deduce

$$\|\zeta_h(t)\|_{0,\Omega}^2 \leq C e^{CT} (\|\zeta_h(0)\|_{0,\Omega}^2 + \|\xi_h(0)\|_{1,\Omega_2}^2 + \|\partial_t \mathbf{f}\|_{L^2(0,T;\mathbf{L}^2(\Omega))}^2).$$

The last estimate and (3.52) yield

$$(3.54) \quad \begin{aligned} & \|\zeta_h(t)\|_{0,\Omega}^2 + \int_0^t a_1[\zeta_h(t), \zeta_h(t)] dt + a_2 \left[\int_0^t \zeta_h(s) ds, \int_0^t \zeta_h(s) ds \right] \\ & \leq C e^{CT} (\|\zeta_h(0)\|_{0,\Omega}^2 + \|\partial_t \mathbf{f}\|_{L^2(0,T;\mathbf{L}^2(\Omega))}^2 + \|\xi_h(0)\|_{1,\Omega_2}^2). \end{aligned}$$

The term $\|\xi_h(0)\|_{1,\Omega_2}^2$ on the right-hand side of (3.54) can be estimated with the help of inverse inequalities (3.13), (3.45), and (3.35) with $r = 1$:

$$(3.55) \quad \begin{aligned} & \|\xi_h(0)\|_{1,\Omega_2} \leq \|\xi_{0,h} - \bar{\mathbf{u}}_{1,h}\|_{1,\Omega_2} + \|\bar{\mathbf{u}}_{1,h} - \mathbf{u}_1\|_{1,\Omega_2} + \|\mathbf{u}_1\|_{1,\Omega_2} \\ & \leq \frac{C}{h} \|\xi_{0,h} - \bar{\mathbf{u}}_{1,h}\|_{0,\Omega_2} + Ch \|\mathbf{u}_1\|_{2,\Omega_2} + \|\mathbf{u}_1\|_{1,\Omega_2} \\ & \leq \frac{C}{h} \|\xi_{0,h} - \mathbf{u}_1\|_{0,\Omega_2} + \frac{C}{h} \|\mathbf{u}_1 - \bar{\mathbf{u}}_{1,h}\|_{0,\Omega_2} + Ch \|\mathbf{u}_1\|_{2,\Omega_2} + \|\mathbf{u}_1\|_{1,\Omega_2} \\ & \leq C(\|\mathbf{v}_0\|_{2,\Omega_1} + \|\mathbf{u}_1\|_{2,\Omega_2} + \|p_0\|_{0,\Omega_1}), \end{aligned}$$

where $\bar{\mathbf{u}}_{1,h}$ is defined by (3.44). The term $\|\boldsymbol{\xi}_h(0)\|_{0,\Omega}^2$ can be estimated as follows. Evaluating (3.21) at $t = 0$, then setting $\boldsymbol{\eta}_h = \partial_t \boldsymbol{\xi}_h(0)$ and using (3.52), we have

$$\begin{aligned}
[[\partial_t \boldsymbol{\xi}_h(0), \partial_t \boldsymbol{\xi}_h(0)]] &= [[\mathbf{f}(0), \partial_t \boldsymbol{\xi}_h(0)]] - a_2[\mathbf{u}_0, \partial_t \boldsymbol{\xi}_h(0)] - a_1[\boldsymbol{\xi}_h(0), \partial_t \boldsymbol{\xi}_h(0)] \\
&= [[\mathbf{f}(0), \partial_t \boldsymbol{\xi}_h(0)]] - a_2[\mathbf{u}_0, \partial_t \boldsymbol{\xi}_h(0)] - b[\partial_t \boldsymbol{\xi}_h(0), p_0] - a_1[\mathbf{v}_0, \partial_t \boldsymbol{\xi}_h(0)] \\
&\quad - [\mathbf{u}_1, \partial_t \boldsymbol{\xi}_h(0)]_{\Omega_2} + [\boldsymbol{\xi}_h(0), \partial_t \boldsymbol{\xi}_h(0)]_{\Omega_2} \\
&= [[\mathbf{f}(0), \partial_t \boldsymbol{\xi}_h(0)]] + [\Delta \mathbf{u}_0 + \nabla(\operatorname{div} \mathbf{u}_0), \partial_t \boldsymbol{\xi}_h(0)]_{\Omega_2} + [\Delta \mathbf{v}_0 - \nabla p_0, \partial_t \boldsymbol{\xi}_h(0)] \\
&\quad + \int_{\Gamma_0} (-\mu_2 \nabla \mathbf{u}_0 \cdot \mathbf{n}_2 - (\lambda_2 + \mu_2)(\operatorname{div} \mathbf{u}_0) \mathbf{n}_2 + p_0 \mathbf{n}_1 - \nabla \mathbf{v}_0 \cdot \mathbf{n}_1) \cdot \partial_t \boldsymbol{\xi}_h(0) \, d\Gamma \\
&\quad - [\mathbf{u}_1, \partial_t \boldsymbol{\xi}_h(0)]_{\Omega_2} + [\boldsymbol{\xi}_h(0), \partial_t \boldsymbol{\xi}_h(0)]_{\Omega_2}.
\end{aligned}$$

Applying assumption (3.47) and initial condition (3.32)–(3.33) to the last relation, we are led to

$$\begin{aligned}
[[\partial_t \boldsymbol{\xi}_h(0), \partial_t \boldsymbol{\xi}_h(0)]] &= [[\mathbf{f}(0), \partial_t \boldsymbol{\xi}_h(0)]] + [\Delta \mathbf{u}_0 + \nabla(\operatorname{div} \mathbf{u}_0), \partial_t \boldsymbol{\xi}_h(0)]_{\Omega_2} \\
&\quad + [\Delta \mathbf{v}_0 - \nabla p_0, \partial_t \boldsymbol{\xi}_h(0)] - [\mathbf{u}_1, \partial_t \boldsymbol{\xi}_h(0)]_{\Omega_2} + [\boldsymbol{\xi}_h(0), \partial_t \boldsymbol{\xi}_h(0)]_{\Omega_2} \\
&\leq C(\|\mathbf{f}(0)\|_{\mathbf{L}^2(\Omega)}^2 + \|\mathbf{u}_0\|_{2,\Omega_2}^2 + \|\mathbf{v}_0\|_2^2 + \|\mathbf{u}_1\|_{2,\Omega_2}^2 + \|p_0\|_{1,\Omega_1}^2) \\
&\quad + C\|\boldsymbol{\xi}_{0,h}\|_{0,\Omega}^2 + \frac{1}{2}[[\partial_t \boldsymbol{\xi}_h(0), \partial_t \boldsymbol{\xi}_h(0)]],
\end{aligned}$$

so that, using (3.55), the last relation simplifies to

$$\|\partial_t \boldsymbol{\xi}_h(0)\|_{0,\Omega}^2 \leq C(\|\mathbf{f}\|_{H^1(0,T;\mathbf{L}^2(\Omega))}^2 + \|\mathbf{u}_0\|_{2,\Omega_2}^2 + \|\mathbf{v}_0\|_{2,\Omega_1}^2 + \|p_0\|_{1,\Omega_1}^2 + \|\mathbf{u}_1\|_{2,\Omega_2}^2).$$

Combining (3.54), (3.55), and the last relation, we obtain (3.50). \square

Remark 2. The particular choice of the initial condition (3.32)–(3.33) played a key role in the estimation of $[[\partial_t \boldsymbol{\xi}_h(0), \partial_t \boldsymbol{\xi}_h(0)]]$.

Using relation (3.20) in reverse, i.e., setting $\mathbf{u}_h = \mathbf{u}_{0,h} + \int_0^t \boldsymbol{\xi}_h(s)|_{\Omega_2} \, ds$ and $\mathbf{v}_h = \boldsymbol{\xi}_h|_{\Omega_1}$, we arrive at the following theorem.

THEOREM 3.9. *Assume that $\mathbf{f}_1, \mathbf{v}_0, \mathbf{f}_2, \mathbf{u}_0$, and \mathbf{u}_1 satisfy (2.6) and (3.46). Assume further that there exists a $p_0 \in H^1(\Omega_1)$ such that (3.47) holds. Then there exists a unique solution $(\mathbf{v}_h, p_h, \mathbf{u}_h) \in C^1([0, T]; X_1^h) \times C([0, T]; Q_1^h) \times C^1([0, T]; X_2^h)$ for (3.14)–(3.19) with the initial conditions $(\mathbf{v}_{0,h}, \mathbf{u}_{1,h})$ defined by (3.32)–(3.33). Moreover, $(\mathbf{v}_h, p_h, \mathbf{u}_h)$ satisfies the estimates*

$$\begin{aligned}
&\|\mathbf{v}_h\|_{L^\infty(0,T;\mathbf{L}^2(\Omega_1))}^2 + \|\partial_t \mathbf{u}_h\|_{L^\infty(0,T;\mathbf{L}^2(\Omega_2))}^2 \\
(3.56) \quad &\quad + \|\mathbf{v}_h\|_{L^2(0,T;\mathbf{H}^1(\Omega_1))}^2 + \|\mathbf{u}_h\|_{L^\infty(0,T;\mathbf{H}^1(\Omega_2))}^2 \\
&\leq C e^{CT} (\|\mathbf{f}\|_{L^2(0,T;\mathbf{L}^2(\Omega))}^2 + \|\mathbf{u}_0\|_{1,\Omega_2}^2 + \|\mathbf{v}_0\|_{1,\Omega_1}^2 + \|p_0\|_{0,\Omega_1}^2 + \|\mathbf{u}_1\|_{0,\Omega_2}^2)
\end{aligned}$$

and

$$\begin{aligned}
&\|\partial_t \mathbf{v}_h\|_{L^\infty(0,T;\mathbf{L}^2(\Omega_1))}^2 + \|\partial_{tt} \mathbf{u}_h\|_{L^\infty(0,T;\mathbf{L}^2(\Omega_2))}^2 \\
(3.57) \quad &\quad + \|\partial_t \mathbf{v}_h\|_{L^2(0,T;\mathbf{H}^1(\Omega_1))}^2 + \|\partial_t \mathbf{u}_h\|_{L^\infty(0,T;\mathbf{H}^1(\Omega_2))}^2 \\
&\leq C e^{CT} (\|\mathbf{f}\|_{H^1(0,T;\mathbf{L}^2(\Omega))}^2 + \|\mathbf{u}_0\|_{2,\Omega_2}^2 + \|\mathbf{v}_0\|_{2,\Omega_1}^2 + \|p_0\|_{1,\Omega_1}^2 + \|\mathbf{u}_1\|_{2,\Omega_2}^2).
\end{aligned}$$

Recall that Theorem 3.6 stated only the existence and uniqueness of a discrete pressure p_h satisfying (3.28), (3.27) and (3.14). By virtue of the strong energy estimates (3.57) and the discrete inf-sup conditions, we now can establish an estimate for p_h .

THEOREM 3.10. *Assume that $\mathbf{f}_1, \mathbf{v}_0, \mathbf{f}_2, \mathbf{u}_0$, and \mathbf{u}_1 satisfy (2.6) and (3.46). Assume further that there exists a $p_0 \in H^1(\Omega_1)$ such that (3.47) holds. Let $(\mathbf{v}_h, p_h, \mathbf{u}_h) \in C^1([0, T]; X_1^h) \times C([0, T]; Q_1^h) \times C^1([0, T]; X_2^h)$ be the solution for (3.14)–(3.19) with the initial conditions $(\mathbf{v}_{0,h}, \mathbf{u}_{1,h})$ defined by (3.32)–(3.33). Then p_h satisfies the estimate*

$$(3.58) \quad \begin{aligned} & \|p_h\|_{L^2(0,T;L^2(\Omega_1))}^2 \\ & \leq C e^{CT} \left(\|\mathbf{f}\|_{H^1(0,T;L^2(\Omega))}^2 + \|\mathbf{u}_0\|_{2,\Omega_2}^2 + \|\mathbf{v}_0\|_{2,\Omega_1}^2 + \|p_0\|_{1,\Omega_1}^2 + \|\mathbf{u}_1\|_{2,\Omega_2}^2 \right). \end{aligned}$$

Proof. We observe that from (3.28) we have

$$\begin{aligned} \|p_h\|_{L^2(0,T;L^2(\Omega_1))} & \leq C \left(\|\partial_t \boldsymbol{\xi}_h\|_{L^2(0,T;L^2(\Omega))}^2 \right. \\ & \quad \left. + \|\mathbf{f}\|_{L^2(0,T;L^2(\Omega))}^2 + \|\boldsymbol{\xi}_h\|_{L^2(0,T;X_1)} + \left\| \int_0^t \boldsymbol{\xi}_h(s) ds \right\|_{L^2(0,T;X_1)} \right). \end{aligned}$$

Thus, (3.58) follows from the last relation and energy estimate (3.50) for $\boldsymbol{\xi}_h$. \square

Remark 3. Note that Theorems 3.8, 3.9, and 3.10 require the specification of an initial pressure p_0 and the initial interface stress condition (3.47). From a physical point of view, these requirements are entirely reasonable.

4. The convergence of finite element solutions and error estimates.

Having proved the existence of finite element solutions $(\mathbf{v}_h, p_h, \mathbf{u}_h)$ for problem (3.14)–(3.19) and (3.32)–(3.33), we now prove the convergence of the finite element solutions and derive error estimates.

4.1. The convergence of finite element solutions. We first consider the convergence of the finite element approximations.

THEOREM 4.1. *Assume that $\mathbf{f}_1, \mathbf{v}_0, \mathbf{f}_2, \mathbf{u}_0$, and \mathbf{u}_1 satisfy (2.6) and (3.46) and that there exists a $p_0 \in H^1(\Omega_1)$ such that (3.47) holds. Let $(\mathbf{v}_h, p_h, \mathbf{u}_h) \in C^1([0, T]; X_1^h) \times C([0, T]; Q_1^h) \times C^1([0, T]; X_2^h)$ be the unique solution of (3.14)–(3.19) with the initial conditions $(\mathbf{v}_{0,h}, \mathbf{u}_{1,h})$ defined by (3.32)–(3.33). Assume further that the finite element meshes are nested, i.e., that the triangulation $\mathcal{T}^{h_2}(\Omega)$ is a refinement of the triangulation $\mathcal{T}^{h_1}(\Omega)$ whenever $h_2 < h_1$. Then, there exists a unique $(\mathbf{v}, p, \mathbf{u})$ such that*

$$(4.1) \quad \begin{cases} \mathbf{v} \in L^\infty(0, T; \mathbf{L}^2(\Omega_1)) \cap L^2(0, T; X_1), \\ \partial_t \mathbf{v} \in L^\infty(0, T; \mathbf{L}^2(\Omega_1)) \cap L^2(0, T; X_1), \quad p \in L^2(0, T; L^2(\Omega_1)), \\ \mathbf{u} \in L^\infty(0, T; X_2), \quad \partial_t \mathbf{u} \in L^\infty(0, T; X_2), \quad \partial_{tt} \mathbf{u} \in L^\infty(0, T; \mathbf{L}^2(\Omega_2)), \end{cases}$$

$$(4.2) \quad \mathbf{v}_h \rightharpoonup \mathbf{v} \quad \text{in } L^2(0, T; X_1), \quad \mathbf{v}_h \overset{*}{\rightharpoonup} \mathbf{v} \quad \text{in } L^\infty(0, T; \mathbf{L}^2(\Omega_1)),$$

$$(4.3) \quad \partial_t \mathbf{v}_h \overset{*}{\rightharpoonup} \partial_t \mathbf{v} \quad \text{in } L^\infty(0, T; \mathbf{L}^2(\Omega_1)), \quad \partial_t \mathbf{v}_h \rightharpoonup \partial_t \mathbf{v} \quad \text{in } L^2(0, T; X_1),$$

$$(4.4) \quad \mathbf{u}_h \overset{*}{\rightharpoonup} \mathbf{u} \quad \text{in } L^\infty(0, T; X_2),$$

$$(4.5) \quad \partial_t \mathbf{u}_h \overset{*}{\rightharpoonup} \partial_t \mathbf{u} \quad \text{in } L^\infty(0, T; \mathbf{L}^2(\Omega_1)), \quad \partial_t \mathbf{u}_h \overset{*}{\rightharpoonup} \partial_t \mathbf{u} \quad \text{in } L^\infty(0, T; X_2),$$

$$(4.6) \quad \partial_{tt} \mathbf{u}_h \overset{*}{\rightharpoonup} \partial_{tt} \mathbf{u} \quad \text{in } L^\infty(0, T; \mathbf{L}^2(\Omega_2)),$$

and

$$(4.7) \quad p^h \rightharpoonup p \quad \text{weakly in } L^2(0, T; L^2(\Omega_1)).$$

Furthermore, $(\mathbf{v}, p, \mathbf{u})$ satisfies (3.3)–(3.6) and the estimates

$$(4.8) \quad \begin{aligned} & \|\mathbf{v}\|_{L^\infty(0, T; \mathbf{L}^2(\Omega_1))}^2 + \|\partial_t \mathbf{u}\|_{L^\infty(0, T; \mathbf{L}^2(\Omega_2))}^2 \\ & + \|\mathbf{v}\|_{L^2(0, T; \mathbf{H}^1(\Omega_1))}^2 + \|\mathbf{u}\|_{L^\infty(0, T; \mathbf{H}^1(\Omega_2))}^2 \\ & \leq C e^{CT} (\|\mathbf{f}\|_{L^2(0, T; \mathbf{L}^2(\Omega))}^2 + \|\mathbf{u}_0\|_{2, \Omega_2}^2 + \|\mathbf{v}_0\|_{2, \Omega_1}^2 + \|p_0\|_{1, \Omega_1}^2 + \|\mathbf{u}_1\|_{2, \Omega_2}^2) \end{aligned}$$

and

$$(4.9) \quad \begin{aligned} & \|\partial_t \mathbf{v}\|_{L^\infty(0, T; \mathbf{L}^2(\Omega_1))}^2 + \|\partial_t \mathbf{v}\|_{L^2(0, T; \mathbf{H}^1(\Omega_1))}^2 + \|p\|_{L^2(0, T; L^2(\Omega_1))}^2 \\ & + \|\partial_{tt} \mathbf{u}(t)\|_{L^\infty(0, T; \mathbf{L}^2(\Omega_2))}^2 + \|\partial_t \mathbf{u}\|_{L^\infty(0, T; \mathbf{H}^1(\Omega_2))}^2 \\ & \leq C e^{CT} (\|\mathbf{f}\|_{H^1(0, T; \mathbf{L}^2(\Omega))}^2 + \|\mathbf{u}_0\|_{2, \Omega_2}^2 + \|\mathbf{v}_0\|_{2, \Omega_1}^2 + \|p_0\|_{1, \Omega_1}^2 + \|\mathbf{u}_1\|_{2, \Omega_2}^2). \end{aligned}$$

Proof. We have that $\{(\mathbf{v}_h, p_h, \mathbf{u}_h)\}$ satisfies the estimates (3.56)–(3.57) and (3.58). Using these estimates, we may extract a subsequence $\{(\mathbf{v}_{h_n}, p_{h_n}, \mathbf{u}_{h_n})\}$ of $\{(\mathbf{v}_h, p_h, \mathbf{u}_h)\}$, with $\{h_n\}$ decreasing to 0 as $n \rightarrow \infty$, such that (4.2)–(4.7) hold for the subsequence $\{(\mathbf{v}_{h_n}, p_{h_n}, \mathbf{u}_{h_n})\}$ for a $(\mathbf{v}, p, \mathbf{u})$ satisfying (4.1).

Equation (3.17) holds for $h = h_n$, and thus, by passing to the limit as $n \rightarrow \infty$ in that equation, we obtain (3.6). Also, $\mathbf{u}(0) = \mathbf{u}_0$ trivially holds.

To prove that $(\mathbf{v}, p, \mathbf{u})$ satisfies (3.3) we begin from (3.14) with $h = h_n$. We arbitrarily fix an integer N and a function $\boldsymbol{\eta} \in C^1([0, T]; X^{h_N})$. For each $n > N$ we obtain from (3.14) and the nesting assumption on the triangulation family $\mathcal{T}^h(\Omega)$ that

$$(4.10) \quad \begin{aligned} & \int_0^T \left(\rho_1 [\partial_t \mathbf{v}_{h_n}, \boldsymbol{\eta}]_{\Omega_1} + a_1 [\mathbf{v}_{h_n}, \boldsymbol{\eta}] + b[\boldsymbol{\eta}, p_{h_n}] + \rho_2 [\partial_{tt} \mathbf{u}_{h_n}, \boldsymbol{\eta}]_{\Omega_2} + a_2 [\mathbf{u}_{h_n}, \boldsymbol{\eta}] \right) dt \\ & = \int_0^T \left(\rho_1 [\mathbf{f}_1, \boldsymbol{\eta}]_{\Omega_1} + \rho_2 [\mathbf{f}_2, \boldsymbol{\eta}]_{\Omega_2} \right) dt. \end{aligned}$$

Passing to the limit as $n \rightarrow \infty$, we find

$$(4.11) \quad \begin{aligned} & \int_0^T \left(\rho_1 [\partial_t \mathbf{v}, \boldsymbol{\eta}]_{\Omega_1} + a_1 [\mathbf{v}, \boldsymbol{\eta}] + b[\boldsymbol{\eta}, p] + \rho_2 [\partial_{tt} \mathbf{u}, \boldsymbol{\eta}]_{\Omega_2} + a_2 [\mathbf{u}, \boldsymbol{\eta}] \right) dt \\ & = \int_0^T \left(\rho_1 [\mathbf{f}_1, \boldsymbol{\eta}]_{\Omega_1} + \rho_2 [\mathbf{f}_2, \boldsymbol{\eta}]_{\Omega_2} \right) dt. \end{aligned}$$

Equality (4.11) then holds for all $\boldsymbol{\eta} \in L^2(0, T; \mathbf{H}_0^1(\Omega))$, as $\bigcup_{n=N}^{\infty} C([0, T]; X^{h_n})$ is dense in $L^2(0, T; \mathbf{H}_0^1(\Omega))$ for the $L^2(0, T; \mathbf{H}_0^1(\Omega))$ norm. Hence,

$$\begin{aligned} & \rho_1[\partial_t \mathbf{v}, \boldsymbol{\eta}]_{\Omega_1} + a_1[\mathbf{v}, \boldsymbol{\eta}] + b[\boldsymbol{\eta}, p] + \rho_2[\partial_{tt} \mathbf{u}, \boldsymbol{\eta}]_{\Omega_2} + a_2[\mathbf{u}, \boldsymbol{\eta}] \\ &= \rho_1[\mathbf{f}_1, \boldsymbol{\eta}]_{\Omega_1} + \rho_2[\mathbf{f}_2, \boldsymbol{\eta}]_{\Omega_2} \quad \forall \boldsymbol{\eta} \in \mathbf{H}_0^1(\Omega), \text{ a.e. } t, \end{aligned}$$

which is precisely (3.3).

From (3.15) we obtain

$$\int_0^T b[\mathbf{v}_{h_n}, q] ds = 0$$

for all $q \in L^2(0, T; Q_1^{h_N})$ and all $n \geq N$. Passing to the limit as $n \rightarrow \infty$ leads us to

$$(4.12) \quad \int_0^T b[\mathbf{v}, q] ds = 0$$

for all $q \in L^2(0, T; Q_1^{h_N})$. Using the denseness (with respect to the $L^2(0, T; L^2(\Omega_1))$ norm) of $\bigcup_{n=N}^{\infty} L^2(0, T; Q_1^{h_n})$ in $L^2(0, T; L^2(\Omega_1))$, we see that (4.12) holds for all $q \in L^2(0, T; L^2(\Omega_1))$. In particular, this implies (3.4).

To verify the initial condition (3.5) we first note that the regularity results (4.1) imply that $\mathbf{v} \in C([0, T]; \mathbf{L}^2(\Omega_1)) \cap C([0, T]; X_1)$, $\mathbf{u} \in C([0, T]; \mathbf{L}^2(\Omega_2)) \cap C([0, T]; X_2)$, and $\partial_t \mathbf{u} \in C([0, T]; \mathbf{L}^2(\Omega_2))$. For each $\boldsymbol{\eta} \in C^1([0, T]; \mathbf{H}_0^1(\Omega))$ with $\boldsymbol{\eta}(T) = \mathbf{0}$ we obtain, from (4.11), by integration by parts that

$$(4.13) \quad \begin{aligned} & \int_0^T \left(-\rho_1[\mathbf{v}, \partial_t \boldsymbol{\eta}]_{\Omega_1} - \rho_2[\partial_t \mathbf{u}, \partial_t \boldsymbol{\eta}]_{\Omega_2} + a_1[\mathbf{v}, \boldsymbol{\eta}] + b[\boldsymbol{\eta}, \hat{p}] + a_2[\mathbf{u}, \boldsymbol{\eta}] \right) dt \\ &= \int_0^T [[\mathbf{f}, \boldsymbol{\eta}]] dt + \rho_1[\mathbf{v}(0), \boldsymbol{\eta}(0)]_{\Omega_1} + \rho_2[\partial_t \mathbf{u}(0), \boldsymbol{\eta}(0)]_{\Omega_2}. \end{aligned}$$

On the other hand, from (4.10), we deduce that for all $\boldsymbol{\eta} \in C^1([0, T]; X^{h_N})$ and all $n > N$,

$$(4.14) \quad \begin{aligned} & \int_0^T \left(-\rho_1[\mathbf{v}_{h_n}, \partial_t \boldsymbol{\eta}]_{\Omega_1} - \rho_2[\partial_t \mathbf{u}_{h_n}, \partial_t \boldsymbol{\eta}]_{\Omega_2} \right. \\ & \quad \left. + a_1[\mathbf{v}_{h_n}, \boldsymbol{\eta}] + b[\boldsymbol{\eta}, p_{h_n}] + a_2[\mathbf{u}_{h_n}, \boldsymbol{\eta}] \right) dt \\ &= \int_0^T [[\mathbf{f}, \boldsymbol{\eta}]] dt + \rho_1[\mathbf{v}_{h_n}(0), \boldsymbol{\eta}(0)]_{\Omega_1} + \rho_2[\partial_t \mathbf{u}_{h_n}(0), \boldsymbol{\eta}(0)]_{\Omega_2}. \end{aligned}$$

Holding N fixed and passing to the limit as $n \rightarrow \infty$ in (4.14) and utilizing (3.35), we arrive at

$$(4.15) \quad \begin{aligned} & \int_0^T \left(-\rho_1[\mathbf{v}, \partial_t \boldsymbol{\eta}]_{\Omega_1} - \rho_2[\partial_t \mathbf{u}, \partial_t \boldsymbol{\eta}]_{\Omega_2} + a_1[\mathbf{v}, \boldsymbol{\eta}] + b[\boldsymbol{\eta}, \hat{p}] + a_2[\mathbf{u}, \boldsymbol{\eta}] \right) dt \\ &= \int_0^T [[\mathbf{f}, \boldsymbol{\eta}]] dt + \rho_1[\mathbf{v}_0, \boldsymbol{\eta}(0)]_{\Omega_1} + \rho_2[\mathbf{u}_1, \boldsymbol{\eta}(0)]_{\Omega_2} \end{aligned}$$

for all $\boldsymbol{\eta} \in C^1([0, T]; X^{h_N})$. Comparing (4.13) and (4.15), we obtain

$$(4.16) \quad \rho_1[\mathbf{v}(0) - \mathbf{v}_0, \boldsymbol{\eta}(0)]_{\Omega_1} + \rho_2[\partial_t \mathbf{u}(0) - \mathbf{u}_1, \boldsymbol{\eta}(0)]_{\Omega_2} = 0$$

for all $\boldsymbol{\eta}(0) \in X^{h_N}$. Since $\bigcup_{n=N}^{\infty} X^{h_n}$ is dense in $\mathbf{L}^2(\Omega)$ for the $\mathbf{L}^2(\Omega)$ norm, we derive

$$\mathbf{v}(0) = \mathbf{v}_0 \quad \text{in } \mathbf{L}^2(\Omega_1) \quad \text{and} \quad \partial_t \mathbf{u}(0) = \mathbf{u}_1 \quad \text{in } \mathbf{L}^2(\Omega_2).$$

To check $\mathbf{u}(0) = \mathbf{u}_0$ we first note that with regularity (4.1) we are justified to write

$$(4.17) \quad \mathbf{u} = \mathbf{u}(0) + \int_0^t \partial_t \mathbf{u}(s) \, ds.$$

From the compact embedding $H^1(0, T; B) \hookrightarrow L^2(0, T; B)$ for any Banach space B and the weak convergence (4.2)–(4.5) we deduce that for a further subsequence h_{n_j} we have

$$\partial_t \mathbf{u}_{h_{n_j}} \rightharpoonup \partial_t \mathbf{u} \quad \text{in } L^2(0, T; \mathbf{L}^2(\Omega_2)) \quad \text{and} \quad \mathbf{u}_{h_{n_j}} \rightarrow \mathbf{u} \quad \text{in } L^2(0, T; \mathbf{L}^2(\Omega_2)),$$

so that, passing to the limit in the relation

$$\mathbf{u}_{h_n} = \mathbf{u}_{0, h_n} + \int_0^t \partial_t \mathbf{u}_{h_n}(s) \, ds$$

and noting that $\|\mathbf{u}_{0, h} - \mathbf{u}_0\|_{0, \Omega_2} \rightarrow 0$ as $h \rightarrow 0$, we obtain

$$(4.18) \quad \mathbf{u} = \mathbf{u}_0 + \int_0^t \partial_t \mathbf{u}(s) \, ds.$$

A comparison of (4.17) and (4.18) yields $\mathbf{u}(0) = \mathbf{u}_0$.

Hence we have verified that $(\mathbf{v}, p, \mathbf{u})$ satisfies (3.1)–(3.6). Of course, (\mathbf{v}, \mathbf{u}) is also a solution for (2.7)–(2.10), so that, by Theorem 2.2, (\mathbf{v}, \mathbf{u}) is the unique solution of (2.7)–(2.10) and estimate (4.8) holds. Then, by Theorem 2.3, we obtain the uniqueness of p . Estimate (4.9) follows from (3.57) and (3.58).

Finally, it follows from the uniqueness of the limit $(\mathbf{v}, p, \mathbf{u})$ that the entire family of finite element solutions $(\mathbf{v}_h, p_h, \mathbf{u}_h)$ satisfies (4.2)–(4.7) as $h \rightarrow 0$. \square

We also have the following strong convergence, the proof of which is contained in that of Theorem 4.1.

COROLLARY 4.2. *Assume that all hypotheses of Theorem 4.1 hold. Then*

$$\mathbf{v}_h \rightarrow \mathbf{v} \quad \text{in } L^2(0, T; \mathbf{L}^2(\Omega_1)), \quad \mathbf{u}_h \rightarrow \mathbf{u} \quad \text{in } L^2(0, T; X_2)$$

and

$$\partial_t \mathbf{u}_h \rightarrow \partial_t \mathbf{u} \quad \text{in } L^2(0, T; \mathbf{L}^2(\Omega_2)).$$

4.2. Error estimates for finite element approximations. We will estimate the error between the continuous solution defined by (3.3)–(3.6) and the finite element solution defined by (3.14)–(3.19) and (3.32)–(3.33). To this end we introduce the weighted $\mathbf{L}^2(\Omega)$ projection operator onto the discretely divergence-free space $\boldsymbol{\Psi}^h$. ($\boldsymbol{\Psi}^h$ is discretely divergence-free in Ω_1 .)

The projection operator $\mathcal{P}^h : \mathbf{L}^2(\Omega) \rightarrow \boldsymbol{\Psi}^h$ with respect to the weighted $\mathbf{L}^2(\Omega)$ inner product is defined as follows: for every $\boldsymbol{\eta} \in \mathbf{L}^2(\Omega)$, $\mathcal{P}^h \boldsymbol{\eta} \in \boldsymbol{\Psi}^h$ is the solution of

$$(4.19) \quad [[\mathcal{P}^h \boldsymbol{\eta}, \mathbf{z}^h]] = [[\boldsymbol{\eta}, \mathbf{z}^h]] \quad \forall \mathbf{z}^h \in \boldsymbol{\Psi}^h.$$

Note that the definition of Ψ^h implies

$$(4.20) \quad b[\mathcal{P}^h \boldsymbol{\eta}, q^h] = 0 \quad \forall q^h \in Q_1^h.$$

We assume that the domains Ω_1 and Ω_2 satisfy the following regularity assumptions.

Hypothesis (H1). The problem

$$(4.21) \quad \begin{cases} (\bar{\mathbf{v}}, \bar{p}) \in \mathbf{H}_0^1(\Omega_1) \times L_0^2(\Omega_1), \\ [\nabla \bar{\mathbf{v}}, \nabla \mathbf{z}]_{\Omega_1} + b[\mathbf{z}, \bar{p}] = [\bar{\mathbf{f}}_1, \mathbf{z}]_{\Omega_1} \quad \forall \mathbf{z} \in \mathbf{H}_0^1(\Omega_1), \\ b[\bar{\mathbf{v}}, q] = 0 \quad \forall q \in L^2(\Omega_1) \end{cases}$$

is $\mathbf{H}^{2-\epsilon_1}$ regular for an $\epsilon_1 \in (0, 1)$; i.e., for every $\bar{\mathbf{f}}_1 \in \mathbf{L}^2(\Omega_1)$, the solution $(\bar{\mathbf{v}}, \bar{p})$ to problem (4.21) belongs to $\mathbf{H}^{2-\epsilon_1}(\Omega_1) \times H^{1-\epsilon_1}(\Omega_1)$, $-\bar{p}\mathbf{n}_1 + (\nabla \bar{\mathbf{v}} + \nabla \bar{\mathbf{v}}^T)\mathbf{n}_1 \in \mathbf{H}^{1/2-\epsilon_1}(\Gamma_0)$, and

$$\|\bar{\mathbf{v}}\|_{\mathbf{H}^{2-\epsilon_1}(\Omega_1)} + \|\bar{p}\|_{H^{1-\epsilon_1}(\Omega_1)} \|-\bar{p}\mathbf{n}_1 + (\nabla \bar{\mathbf{v}} + \nabla \bar{\mathbf{v}}^T)\mathbf{n}_1\|_{1/2-\epsilon_1, \Gamma_0} \leq C \|\bar{\mathbf{f}}_1\|_{0, \Omega_1}.$$

Hypothesis (H2). The problem

$$(4.22) \quad \begin{cases} \bar{\mathbf{u}} \in \mathbf{H}_0^1(\Omega_1), \\ [\nabla \bar{\mathbf{u}}, \nabla \mathbf{w}]_{\Omega_1} = [\bar{\mathbf{f}}_2, \mathbf{w}]_{\Omega_1} \quad \forall \mathbf{w} \in \mathbf{H}_0^1(\Omega_2) \end{cases}$$

is $\mathbf{H}^{2-\epsilon_2}$ regular for an $\epsilon_2 \in (0, 1)$; i.e., for every $\bar{\mathbf{f}}_2 \in \mathbf{L}^2(\Omega_2)$, the solution $\bar{\mathbf{u}}$ to problem (4.22) belongs to $\mathbf{H}^{2-\epsilon_2}(\Omega_2)$, $\nabla \bar{\mathbf{u}} \cdot \mathbf{n}_2 \in \mathbf{H}^{1/2-\epsilon_2}(\Gamma_0)$, and

$$\|\bar{\mathbf{u}}\|_{\mathbf{H}^{2-\epsilon_2}(\Omega_2)} \|\nabla \bar{\mathbf{u}} \cdot \mathbf{n}_2\|_{1/2-\epsilon_2, \Gamma_0} \leq C \|\bar{\mathbf{f}}_2\|_{0, \Omega_2}.$$

Remark 4. Hypotheses (H1)–(H2) are simply equivalent to angle conditions on Ω_1 and Ω_2 owing to the well-known regularity results on polygonal domains for boundary value problems (4.21) and (4.22); see [24] and [19]. In particular, if both Ω_1 and Ω_2 are convex (in which case Γ_0 is necessarily a straight line), then ϵ_1 and ϵ_2 can be chosen arbitrarily small.

Under Hypotheses (H1)–(H2), we may prove the following error estimates for the projection operator \mathcal{P}^h :

$$(4.23) \quad \begin{aligned} \|\zeta - \mathcal{P}^h \zeta\|_{1, \Omega} &\leq Ch^{r-\epsilon} (\|\zeta\|_{r+1, \Omega_1} + \|\zeta\|_{r+1, \Omega_2}) \\ \forall \zeta \in \Psi \quad \text{with } \zeta|_{\Omega_i} &\in \mathbf{H}^{r+1}(\Omega_i), \quad i = 1, 2, r \in [0, k], \end{aligned}$$

and

$$(4.24) \quad \begin{aligned} \|\zeta - \mathcal{P}^h \zeta\|_{0, \Omega} &\leq Ch^{r+1-\epsilon} (\|\zeta\|_{r+1, \Omega_1} + \|\zeta\|_{r+1, \Omega_2}) \\ \forall \zeta \in \Psi \quad \text{with } \zeta|_{\Omega_i} &\in \mathbf{H}^{r+1}(\Omega_i), \quad i = 1, 2, r \in [0, k]. \end{aligned}$$

The proof of (4.23)–(4.24) will be given in the appendix, Theorem A.3.

Now we prove the following error estimates for the semidiscrete finite element approximations of the fluid-solid interaction problem.

THEOREM 4.3. *Assume that $\mathbf{f}_1, \mathbf{v}_0, \mathbf{f}_2, \mathbf{u}_0$, and \mathbf{u}_1 satisfy (2.6) and (3.46) and that there exists a $p_0 \in H^1(\Omega_1)$ such that (3.47) holds. Assume also that (H1)–(H2) hold.*

Let $(\mathbf{v}, p, \mathbf{u})$ be the solution of (3.1)–(3.6), and $(\mathbf{v}_h, p_h, \mathbf{u}_h)$ be the solution of (3.14)–(3.19) and (3.32)–(3.33). Assume that for some $r \in [1, k]$, $\mathbf{v} \in L^2(0, T; \mathbf{H}^{r+1}(\Omega_1))$, $\partial_t \mathbf{v} \in L^2(0, T; \mathbf{H}^{r-1}(\Omega_1))$, $p \in L^2(0, T; H^r(\Omega_1))$, $\partial_t \mathbf{u} \in L^2(0, T; \mathbf{H}^{r+1}(\Omega_2))$, $\partial_{tt} \mathbf{u} \in L^2(0, T; \mathbf{H}^{r-1}(\Omega_2))$, $\mathbf{v}_0 \in \mathbf{H}^{r+1}(\Omega_1)$, $\mathbf{u}_1 \in \mathbf{H}^{r+1}(\Omega_2)$, $\mathbf{u}_0 \in \mathbf{H}^{r+1}(\Omega_2)$, and $p_0 \in H^r(\Omega_1)$. Then,

$$\begin{aligned}
& \|\mathbf{v}(t) - \mathbf{v}_h(t)\|_{0, \Omega_1}^2 + \|\mathbf{v} - \mathbf{v}_h\|_{L^2(0, T; X_1)}^2 \\
& \quad + \|\partial_t \mathbf{u}(t) - \partial_t \mathbf{u}_h(t)\|_{0, \Omega_2}^2 + \|\mathbf{u}(t) - \mathbf{u}_h(t)\|_{1, \Omega_2}^2 \\
(4.25) \quad & \leq C e^{CT} h^{2r} (\|\mathbf{v}_0\|_{r+1, \Omega_1}^2 + \|\mathbf{u}_1\|_{r+1, \Omega_2}^2 + \|\mathbf{u}_0\|_{r+1, \Omega_2}^2 + \|p_0\|_{r, \Omega_1}^2 \\
& \quad + \|p\|_{L^2(0, T; H^r(\Omega_1))}^2) + C e^{CT} h^{2(r-\epsilon)} (\|\mathbf{v}\|_{L^2(0, T; \mathbf{H}^{r+1}(\Omega_1))}^2 \\
& \quad + \|\mathbf{u}_t\|_{L^2(0, T; \mathbf{H}^{r+1}(\Omega_2))}^2 + \|\partial_t \mathbf{v}\|_{L^2(0, T; \mathbf{H}^{r-1}(\Omega_1))}^2 + \|\partial_{tt} \mathbf{u}\|_{L^2(0, T; \mathbf{H}^{r-1}(\Omega_2))}^2)
\end{aligned}$$

for all $t \in [0, T]$.

Proof. Let $\boldsymbol{\xi}$ and $\boldsymbol{\xi}_h$ be defined by (2.11) and (3.20), respectively. We set $\tilde{\mathbf{v}}_h(t) = [\mathcal{P}^h \boldsymbol{\xi}(t)]_{\Omega_1}$ and $\tilde{\mathbf{w}}_h(t) = [\mathcal{P}^h \boldsymbol{\xi}(t)]_{\Omega_2}$.

By subtracting (3.14)–(3.15) from the corresponding equations of (3.3)–(3.4), we obtain the following “orthogonality conditions”:

$$\begin{aligned}
(4.26) \quad & \rho_1 [\partial_t \mathbf{v} - \partial_t \mathbf{v}_h, \boldsymbol{\eta}_h]_{\Omega_1} + b[\boldsymbol{\eta}_h, p - p_h] + a_1 [\mathbf{v} - \mathbf{v}_h, \boldsymbol{\eta}_h] \\
& \quad + \rho_2 [\mathbf{u}_{tt} - \partial_{tt} \mathbf{u}_h, \boldsymbol{\eta}_h]_{\Omega_2} + a_2 [\mathbf{u} - \mathbf{u}_h, \boldsymbol{\eta}_h] = 0 \quad \forall \boldsymbol{\eta}_h \in X^h, \text{ a.e. } t,
\end{aligned}$$

$$(4.27) \quad b[\mathbf{v} - \mathbf{v}_h, q_h] = 0 \quad \forall q_h \in Q_1^h, \text{ a.e. } t.$$

By adding/subtracting terms and using (4.26)–(4.27), we deduce that

$$\begin{aligned}
(4.28) \quad & \rho_1 [\partial_t \mathbf{v}_h - \partial_t \mathbf{v}_h, \mathbf{v} - \mathbf{v}_h]_{\Omega_1} + a_1 [\mathbf{v} - \mathbf{v}_h, \mathbf{v} - \mathbf{v}_h] \\
& \quad + \rho_2 [\partial_{tt} \mathbf{u} - \partial_{tt} \mathbf{u}_h, \partial_t \mathbf{u} - \partial_t \mathbf{u}_h]_{\Omega_2} + a_2 [\mathbf{u} - \mathbf{u}_h, \partial_t \mathbf{u} - \mathbf{u}_h] \\
& = \rho_1 [\partial_t \mathbf{v} - \partial_t \mathbf{v}_h, \mathbf{v} - \tilde{\mathbf{v}}_h]_{\Omega_1} + a_1 [\mathbf{v} - \mathbf{v}_h, \mathbf{v} - \tilde{\mathbf{v}}_h] \\
& \quad + \rho_2 [\partial_{tt} \mathbf{u} - \partial_{tt} \mathbf{u}_h, \partial_t \mathbf{u} - \tilde{\mathbf{w}}_h]_{\Omega_2} + a_2 [\mathbf{u} - \mathbf{u}_h, \partial_t \mathbf{u} - \tilde{\mathbf{w}}_h] \\
& \quad - b[\tilde{\mathbf{v}}_h - \mathbf{v}_h, p - p_h] + \rho_1 [\partial_t \mathbf{v} - \partial_t \mathbf{v}_h, \tilde{\mathbf{v}}_h - \mathbf{v}_h]_{\Omega_1} \\
& \quad + a_1 [\mathbf{v} - \mathbf{v}_h, \tilde{\mathbf{v}} - \mathbf{v}_h] + \rho_2 [\partial_{tt} \mathbf{u} - \partial_{tt} \mathbf{u}_h, \tilde{\mathbf{w}}_h - \partial_t \mathbf{u}_h]_{\Omega_2} \\
& \quad + a_2 [\mathbf{u} - \mathbf{u}_h, \tilde{\mathbf{w}}_h - \partial_t \mathbf{u}_h] + b[\tilde{\mathbf{v}}_h - \mathbf{v}_h, p - p_h] \\
& = \rho_1 [\partial_t \mathbf{v} - \partial_t \mathbf{v}_h, \mathbf{v} - \tilde{\mathbf{v}}_h]_{\Omega_1} + a_1 [\mathbf{v} - \mathbf{v}_h, \mathbf{v} - \tilde{\mathbf{v}}_h] \\
& \quad + \rho_2 [\partial_{tt} \mathbf{u} - \partial_{tt} \mathbf{u}_h, \partial_t \mathbf{u} - \tilde{\mathbf{w}}_h]_{\Omega_2} + a_2 [\mathbf{u} - \mathbf{u}_h, \partial_t \mathbf{u} - \tilde{\mathbf{w}}_h] \\
& \quad + b[\mathbf{v}_h - \tilde{\mathbf{v}}_h, p - p_h].
\end{aligned}$$

By the definition of $\tilde{\mathbf{v}}_h$ and (4.20), we obtain

$$(4.29) \quad b[\tilde{\mathbf{v}}_h(t), p_h] = b[\mathcal{P}^h \boldsymbol{\xi}(t), p_h] = 0 = b[\mathcal{P}^h \boldsymbol{\xi}(t), q_h] = b[\tilde{\mathbf{v}}_h(t), q_h] \quad \forall q_h \in Q_1^h.$$

Utilizing (3.15), we have

$$(4.30) \quad b[\mathbf{v}_h(t), p_h] = 0 = b[\mathbf{v}_h(t), q_h] \quad \forall q_h \in Q_1^h.$$

Additionally,

$$(4.31) \quad \begin{aligned} & \rho_1[\partial_t \mathbf{v} - \partial_t \mathbf{v}_h, \mathbf{v} - \tilde{\mathbf{v}}_h]_{\Omega_1} + \rho_2[\partial_{tt} \mathbf{u} - \partial_{tt} \mathbf{u}_h, \partial_t \mathbf{u} - \tilde{\mathbf{w}}]_{\Omega_2} \\ &= [[\partial_t \boldsymbol{\xi}(t) - \partial_t \boldsymbol{\xi}_h(t), \boldsymbol{\xi}(t) - \mathcal{P}^h \boldsymbol{\xi}(t)]] = [[\partial_t \boldsymbol{\xi}(t), \boldsymbol{\xi}(t) - \mathcal{P}^h \boldsymbol{\xi}(t)]] \\ &= [[\partial_t \boldsymbol{\xi}(t) - \partial_t \mathcal{P}^h \boldsymbol{\xi}(t), \boldsymbol{\xi}(t) - \mathcal{P}^h \boldsymbol{\xi}(t)]] \\ &= \frac{1}{2} \frac{d}{dt} [[\boldsymbol{\xi}(t) - \mathcal{P}^h \boldsymbol{\xi}(t), \boldsymbol{\xi}(t) - \mathcal{P}^h \boldsymbol{\xi}(t)]] \\ &= \frac{\rho_1}{2} \frac{d}{dt} \|\mathbf{v} - \tilde{\mathbf{v}}_h\|_{0, \Omega_1}^2 + \frac{\rho_2}{2} \frac{d}{dt} \|\partial_t \mathbf{u} - \tilde{\mathbf{w}}_h\|_{0, \Omega_2}^2. \end{aligned}$$

Combining (4.28)–(4.31), we deduce that for all $q_h \in L^2(0, T; Q_1^h)$

$$\begin{aligned} & \frac{\rho_1}{2} \frac{d}{dt} \|\mathbf{v} - \mathbf{v}_h\|_{0, \Omega_1}^2 + a_1[\mathbf{v} - \mathbf{v}_h, \mathbf{v} - \mathbf{v}_h] + \frac{\rho_2}{2} \frac{d}{dt} \|\partial_t \mathbf{u} - \partial_t \mathbf{u}_h\|_{0, \Omega_2}^2 \\ & \quad + \frac{1}{2} \frac{d}{dt} a_2[\mathbf{u} - \mathbf{u}_h, \mathbf{u} - \mathbf{u}_h] \\ &= \frac{\rho_1}{2} \frac{d}{dt} \|\mathbf{v} - \tilde{\mathbf{v}}_h\|_{0, \Omega_1}^2 + a_1[\mathbf{v} - \mathbf{v}_h, \mathbf{v} - \tilde{\mathbf{v}}_h] + \frac{\rho_2}{2} \frac{d}{dt} \|\partial_t \mathbf{u} - \tilde{\mathbf{w}}_h\|_{0, \Omega_2}^2 \\ & \quad + a_2[\mathbf{u} - \mathbf{u}_h, \partial_t \mathbf{u} - \tilde{\mathbf{w}}_h] + b[\mathbf{v}_h - \tilde{\mathbf{v}}_h, p - q_h] \\ &\leq \frac{\rho_1}{2} \frac{d}{dt} \|\mathbf{v} - \tilde{\mathbf{v}}_h\|_{0, \Omega_1}^2 + \frac{k_1}{4} \|\mathbf{v}(t) - \mathbf{v}_h(t)\|_{1, \Omega_1}^2 + C \|\mathbf{v}(t) - \tilde{\mathbf{v}}_h(t)\|_{1, \Omega_1}^2 \\ & \quad + \frac{\rho_2}{2} \frac{d}{dt} \|\partial_t \mathbf{u} - \tilde{\mathbf{w}}_h\|_{0, \Omega_2}^2 + \|\mathbf{u}(t) - \mathbf{u}_h(t)\|_{0, \Omega_2}^2 + C \|\partial_t \mathbf{u}(t) - \tilde{\mathbf{w}}_h(t)\|_{1, \Omega_2}^2 \\ & \quad + C \|\mathbf{v}(t) - \tilde{\mathbf{v}}_h(t)\|_{1, \Omega_2}^2 + \frac{k_1}{4} \|\mathbf{v}(t) - \mathbf{v}_h(t)\|_{1, \Omega_2}^2 + C \|p(t) - q_h\|_{0, \Omega_1}^2. \end{aligned}$$

Applying (2.2)–(2.3) to the last relation and integrating in t , we obtain

$$(4.32) \quad \begin{aligned} & \rho_1 \|\mathbf{v}(t) - \mathbf{v}_h(t)\|_{0, \Omega_1}^2 + k_1 \|\mathbf{v} - \mathbf{v}_h\|_{L^2(0, T; \mathbf{H}^1(\Omega_1))}^2 \\ & \quad + \rho_2 \|\partial_t \mathbf{u}(t) - \partial_t \mathbf{u}_h(t)\|_{0, \Omega_2}^2 + \|\mathbf{u}(t) - \mathbf{u}_h(t)\|_{1, \Omega_2}^2 \\ &\leq C \left(\|\mathbf{v}(0) - \mathbf{v}_{0, h}\|_{0, \Omega_1}^2 + \|\partial_t \mathbf{u}(0) - \mathbf{u}_{1, h}\|_{0, \Omega_2}^2 + \|\mathbf{u}_0 - \mathbf{u}_{0, h}\|_{1, \Omega_2}^2 \right. \\ & \quad \left. + \|\boldsymbol{\xi}_0 - \mathcal{P}^h \boldsymbol{\xi}_0\|_{0, \Omega}^2 + \|\boldsymbol{\xi}(t_0) - \mathcal{P}^h \boldsymbol{\xi}(t_0)\|_{0, \Omega_1}^2 + \|\boldsymbol{\xi} - \mathcal{P}^h \boldsymbol{\xi}\|_{L^2(0, T; \mathbf{H}^1(\Omega))}^2 \right. \\ & \quad \left. + \|p - q_h\|_{L^2(0, T; L^2(\Omega_1))}^2 \right) + \int_0^t \|\mathbf{u}(s) - \mathbf{u}_h(s)\|_{1, \Omega_2}^2 ds \end{aligned}$$

for all $q_h \in L^2(0, T; Q_1^h)$, where $t_0 \in [0, T]$ is such that

$$\|\boldsymbol{\xi}(t_0) - \mathcal{P}^h \boldsymbol{\xi}(t_0)\|_{0, \Omega}^2 = \max_{t \in [0, T]} \|\boldsymbol{\xi}(t) - \mathcal{P}^h \boldsymbol{\xi}(t)\|_{0, \Omega}^2.$$

The error estimate (3.34) yields

$$(4.33) \quad \begin{aligned} & \|\mathbf{v}_0 - \mathbf{v}_{0,h}\|_{0,\Omega_1}^2 + \|\mathbf{u}_1 - \mathbf{u}_{1,h}\|_{0,\Omega_2}^2 \\ &= Ch^{2r} (\|\mathbf{v}_0\|_{r+1,\Omega_1}^2 + \|\mathbf{u}_1\|_{r+1,\Omega_2}^2 + \|p_0\|_{r,\Omega_1}^2). \end{aligned}$$

Equation (3.19) and the approximation properties imply

$$(4.34) \quad \|\mathbf{u}_0 - \mathbf{u}_{0,h}\|_{1,\Omega_2}^2 \leq Ch^{2r} \|\mathbf{u}_0\|_{r+1,\Omega_1}^2.$$

Also, by virtue of (4.24), we have

$$(4.35) \quad \begin{aligned} & \|\boldsymbol{\xi}(t_0) - \mathcal{P}^h \boldsymbol{\xi}(t_0)\|_{0,\Omega}^2 \leq Ch^{2r-2\epsilon} \left(\|\mathbf{v}(t_0)\|_{r,\Omega_1}^2 + \|\partial_t \mathbf{u}(t_0)\|_{r,\Omega_2}^2 \right) \\ & \leq Ch^{2r-2\epsilon} \left(\|\mathbf{v}\|_{L^2(0,T;\mathbf{H}^{r+1}(\Omega_1))}^2 + \|\partial_t \mathbf{v}\|_{L^2(0,T;\mathbf{H}^{r-1}(\Omega_1))}^2 \right. \\ & \quad \left. + \|\mathbf{u}\|_{L^2(0,T;\mathbf{H}^{r+1}(\Omega_2))}^2 + \|\partial_t \mathbf{u}\|_{L^2(0,T;\mathbf{H}^{r-1}(\Omega_2))}^2 \right). \end{aligned}$$

Thus, utilizing (4.33)–(4.35), (4.23), and (3.9), we may simplify (4.32) to

$$(4.36) \quad \begin{aligned} & \rho_1 \|\mathbf{v}(t) - \mathbf{v}_h(t)\|_{0,\Omega_1}^2 + k_1 \|\mathbf{v} - \mathbf{v}_h\|_{L^2(0,T;\mathbf{H}^1(\Omega_1))}^2 \\ & \quad + \rho_2 \|\partial_t \mathbf{u}(t) - \partial_t \mathbf{u}_h(t)\|_{0,\Omega_2}^2 + \|\mathbf{u}(t) - \mathbf{u}_h(t)\|_{1,\Omega_2}^2 \\ & \leq Ch^{2r} \left(\|\mathbf{v}_0\|_{r+1,\Omega_1}^2 + \|\mathbf{u}_1\|_{r+1,\Omega_2}^2 + \|p_0\|_{r,\Omega_1}^2 + \|\mathbf{u}_0\|_{r+1,\Omega_1}^2 \right. \\ & \quad \left. + \|p\|_{L^2(0,T;H^r(\Omega_1))}^2 \right) + Ch^{2r-2\epsilon} \left(\|\mathbf{v}\|_{L^2(0,T;\mathbf{H}^{r+1}(\Omega_1))}^2 \right. \\ & \quad \left. + \|\partial_t \mathbf{v}\|_{L^2(0,T;\mathbf{H}^{r-1}(\Omega_1))}^2 + \|\mathbf{u}\|_{L^2(0,T;\mathbf{H}^{r+1}(\Omega_2))}^2 \right. \\ & \quad \left. + \|\partial_t \mathbf{u}\|_{L^2(0,T;\mathbf{H}^{r-1}(\Omega_2))}^2 \right) + \int_0^t \|\mathbf{u}(s) - \mathbf{u}_h(s)\|_{1,\Omega_2}^2 ds. \end{aligned}$$

By dropping the first three terms on the left-hand side of (4.36) and applying the Gronwall's inequality (3.53), we obtain

$$(4.37) \quad \begin{aligned} & \|\mathbf{u}(t) - \mathbf{u}_h(t)\|_{1,\Omega_2}^2 \leq Ce^{CT} h^{2r} \left[\|\mathbf{v}_0\|_{r+1,\Omega_1}^2 + \|\mathbf{u}_1\|_{r+1,\Omega_2}^2 \right. \\ & \quad \left. + \|p_0\|_{r,\Omega_1}^2 + \|\mathbf{u}_0\|_{r+1,\Omega_1}^2 + \|p\|_{L^2(0,T;H^r(\Omega_1))}^2 \right] \\ & \quad + Ce^{CT} h^{2r-2\epsilon} \left(\|\mathbf{v}\|_{L^2(0,T;\mathbf{H}^{r+1}(\Omega_1))}^2 + \|\partial_t \mathbf{v}\|_{L^2(0,T;\mathbf{H}^{r-1}(\Omega_1))}^2 \right. \\ & \quad \left. + \|\mathbf{u}\|_{L^2(0,T;\mathbf{H}^{r+1}(\Omega_2))}^2 + \|\partial_t \mathbf{u}\|_{L^2(0,T;\mathbf{H}^{r-1}(\Omega_2))}^2 \right). \end{aligned}$$

Hence, (4.25) follows from (4.36)–(4.37). \square

Appendix. Error estimates for the weighted L^2 projection onto Ψ^h . The objective of this subsection is to prove error estimates (4.23)–(4.24) for the weighted L^2 projection operator \mathcal{P}^h defined by (4.19).

We introduce an operator $\mathcal{S}^h : \Psi \rightarrow \Psi^h$ as follows. For each $\boldsymbol{\zeta} \in \Psi \subset \mathbf{H}_0^1(\Omega)$,

$$(A.1) \quad \mathcal{S}^h \boldsymbol{\zeta} = \begin{cases} \boldsymbol{\zeta}_{1,h} & \text{in } \Omega_1, \\ \boldsymbol{\zeta}_{2,h} & \text{in } \Omega_2, \end{cases}$$

where $\zeta_{1,h} \in X_1^h$ together with some $\sigma_h \in Q_1^h$ is the finite element solution of

$$\begin{cases} a_1[\zeta_{1,h}, \mathbf{z}_h] + b[\mathbf{z}_h, \bar{\sigma}_h] = [\zeta, \mathbf{z}_h] & \forall \mathbf{z}_h \in X_1^h \cap \mathbf{H}_0^1(\Omega_1), \\ b[\zeta_{1,h}, q_h] = 0 & \forall q_h \in Q_1^h \cap L_0^2(\Omega_1), \\ \zeta_{1,h}|_{\Gamma_1} = \mathbf{0} & \text{and} \quad [\zeta_{1,h}, \mathbf{s}_h]_{0,\Gamma_0} = [\zeta, \mathbf{s}_h]_{0,\Gamma_0} \quad \forall \mathbf{s}_h \in X_1^h|_{\Gamma_0}, \end{cases}$$

and $\zeta_{2,h} \in X_2^h$ is the finite element solution of

$$\begin{cases} [\nabla \zeta_{2,h}, \nabla \mathbf{w}_h]_{\Omega_2} = [\nabla \zeta, \nabla \mathbf{w}_h]_{\Omega_2} & \forall \mathbf{w}_h \in X_2^h \cap \mathbf{H}_0^1(\Omega_2), \\ \zeta_{2,h}|_{\Gamma_2} = \mathbf{0} & \text{and} \quad [\zeta_{2,h}, \mathbf{s}_h]_{0,\Gamma_0} = [\zeta, \mathbf{s}_h]_{0,\Gamma_0} \quad \forall \mathbf{s}_h \in X_2^h|_{\Gamma_0}. \end{cases}$$

Evidently, $\zeta_{1,h}|_{\Gamma_0} = \zeta_{2,h}|_{\Gamma_0}$, so that $\mathcal{S}^h \zeta$ defined by (A.1) indeed satisfies $\mathcal{S}^h \zeta \in \Psi^h$.

Using the results of [23, 25] concerning error estimates for the finite element approximations of the Stokes equations (noting that $\operatorname{div} \zeta|_{\Omega_1} = 0$) with inhomogeneous boundary conditions, we obtain

$$(A.2) \quad \|\zeta_{1,h} - \zeta\|_{1,\Omega_1} \leq Ch^r \|\zeta\|_{r+1,\Omega_1} \quad \text{if } \zeta|_{\Omega_1} \in \mathbf{H}^{r+1}(\Omega_1).$$

Furthermore, under assumption (H1), we may adapt straightforwardly the proof in [23] for an Aubin–Nitsche-type result to obtain

$$(A.3) \quad \|\zeta_{1,h} - \zeta\|_{0,\Omega_1} \leq Ch^{1-\epsilon_1} \|\zeta_{1,h} - \zeta\|_{1,\Omega_1}.$$

Likewise,

$$(A.4) \quad \|\zeta_{2,h} - \zeta\|_{1,\Omega_2} \leq Ch^r \|\zeta\|_{r+1,\Omega_2} \quad \text{if } \zeta|_{\Omega_2} \in \mathbf{H}^{r+1}(\Omega_2),$$

and, under assumption (H2),

$$(A.5) \quad \|\zeta_{2,h} - \zeta\|_{0,\Omega_2} \leq Ch^{1-\epsilon_2} \|\zeta_{2,h} - \zeta\|_{1,\Omega_2}.$$

To summarize, we have the following results.

PROPOSITION A.1. *If $\zeta \in \Psi$ and $\zeta|_{\Omega_i} \in \mathbf{H}^{r+1}(\Omega_i)$ ($i = 1, 2$) for some $r \in [0, k]$, then*

$$(A.6) \quad \|\mathcal{S}^h \zeta - \zeta\|_{1,\Omega} \leq Ch^r (\|\zeta\|_{r+1,\Omega_1} + \|\zeta\|_{r+1,\Omega_2}).$$

If, in addition, assumptions (H1)–(H2) hold, then

$$(A.7) \quad \|\mathcal{S}^h \zeta - \zeta\|_{0,\Omega} \leq Ch^{1-\epsilon} \|\mathcal{S}^h \zeta - \zeta\|_{1,\Omega},$$

where $\epsilon = \max\{\epsilon_1, \epsilon_2\}$

The following proposition establishes relationships between approximation properties for the operator \mathcal{P}^h and those for the operator \mathcal{S}^h .

PROPOSITION A.2. *Assume that (H1)–(H2) hold. Then,*

$$(A.8) \quad \|\zeta - \mathcal{P}^h \zeta\|_{1,\Omega} \leq Ch^{-\epsilon} \|\zeta - \mathcal{S}^h \zeta\|_{1,\Omega} \quad \forall \zeta \in \Psi.$$

Proof. Let $\zeta \in \Psi$ be given. The best approximation property of a projection operator implies that

$$(A.9) \quad \|\zeta - \mathcal{P}^h \zeta\|_{0,\Omega} \leq \|\mathcal{S}^h \zeta - \zeta\|_{0,\Omega}.$$

Using the triangle inequality, the inverse inequality (3.13), and inequality (A.9), we deduce that

$$\begin{aligned}
\|\zeta - \mathcal{P}^h \zeta\|_{1,\Omega} &\leq \|\zeta - \mathcal{S}^h \zeta\|_{1,\Omega} + \|\mathcal{S}^h \zeta - \mathcal{P}^h \zeta\|_{1,\Omega} \\
&\leq \|\zeta - \mathcal{S}^h \zeta\|_{1,\Omega} + \frac{C}{h} \|\mathcal{S}^h \zeta - \mathcal{P}^h \zeta\|_{0,\Omega} \\
&\leq \|\zeta - \mathcal{S}^h \zeta\|_{1,\Omega} + \frac{C}{h} \|\zeta - \mathcal{P}^h \zeta\|_{0,\Omega} + \frac{C}{h} \|\mathcal{S}^h \zeta - \zeta\|_{0,\Omega} \\
&\leq \|\zeta - \mathcal{S}^h \zeta\|_{1,\Omega} + \frac{C}{h} \|\mathcal{S}^h \zeta - \zeta\|_{0,\Omega}.
\end{aligned}$$

Thus, (A.8) follows from the last inequality and (A.7). \square

Finally, as obvious consequences of (A.8) and (A.6)–(A.7), we obtain the following error estimates for $\zeta - \mathcal{P}^h \zeta$:

THEOREM A.3. *Assume that (H1)–(H2) hold. Then the operator \mathcal{P}^h satisfies the error estimates (4.23) and (4.24).*

REFERENCES

- [1] M. BERGGREN, *Approximations of Very Weak Solutions to Boundary-Value Problems*, SIAM J. Numer. Anal., to appear.
- [2] A. BERMÚDEZ, R. DURÁN, AND R. RODRÍGUEZ, *Finite element analysis of compressible and incompressible fluid-solid systems*, Math. Comp., 67 (1998), pp. 111–136.
- [3] F. BLOM, *A monolithic fluid-structure interaction algorithm applied to the piston problem*, Comput. Methods Appl. Mech. Engrg., 167 (1998), pp. 369–391.
- [4] J. BOUJOT, *Mathematical formulation of fluid-structure interaction problems*, RAIRO Modél. Math. Anal. Numér., 21 (1987), pp. 239–260.
- [5] P. CIARLET, *The Finite Element Method For Elliptic Problems*, North-Holland, Amsterdam, 1978.
- [6] C. CONCA AND M. DURÁN, *A numerical study of a spectral problem in solid-fluid type structures* Numer. Methods Partial Differential Equations, 11 (1995), pp. 423–444.
- [7] C. CONCA, J. MARTÍN, AND J. TUCSNAK, *Motion of a rigid body in a viscous fluid*, C. R. Acad. Sci. Paris Sér. I Math., 328 (1999), pp. 473–478.
- [8] S. DASSER, *A penalization method for the homogenization of a mixed fluid-structure problem*, C. R. Acad. Sci. Paris Sér. I Math., 320 (1995), pp. 759–764.
- [9] B. DESJARDINS AND M. J. ESTEBAN, *On weak solutions for fluid-rigid structure interaction: Compressible and incompressible models*, Comm. Partial Differential Equations, 25 (2000), pp. 1399–1413.
- [10] Q. DU, M. GUNZBURGER, L. HOU, AND J. LEE, *Analysis of a Linear Fluid-Structure Interaction Problem*, Disc. Cont. Dyn. Syst., 9 (2003), pp. 633–650.
- [11] D. ERRATE, M. J. ESTEBAN, AND Y. MADAY, *Couplage fluid-structure, Un modèle simplifié en dimension 1*, C. R. Acad. Sci. Paris Sér. I Math., 318 (1994), pp. 275–281.
- [12] L. C. EVANS, *Partial Differential Equations*, AMS, Providence, RI, 1998.
- [13] C. FARHAT, M. LESOINNE, AND P. LETALLEC, *Load and motion transfer algorithms for fluid/structure interaction problems with non-matching discrete interfaces: Momentum and energy conservation, optimal discretization and application to aeroelasticity*, Comput. Methods Appl. Mech. Engrg., 157 (1998), pp. 95–114.
- [14] F. FLORI AND P. ORENGA, *On a fluid-structure interaction problem*, in Trends in Applications of Mathematics to Mechanics (Nice, 1998), Chapman & Hall/CRC, Boca Raton, FL, 2000, pp. 293–305.
- [15] F. FLORI AND P. ORENGA, *Analysis of a nonlinear fluid-structure interaction problem in velocity-displacement formulation*, Nonlinear Anal., 35 (1999), pp. 561–587.
- [16] F. FLORI AND P. ORENGA, *On a nonlinear fluid-structure interaction problem defined on a domain depending on time*, Nonlinear Anal., 38 (1999), pp. 549–569.
- [17] F. FLORI AND P. ORENGA, *Fluid-structure interaction: Analysis of a 3-D compressible model*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 17 (2000), pp. 753–777.
- [18] L. GASTALDI, *Mixed finite element methods in fluid structure system*, Numer. Math., 74 (1996), pp. 153–176.

- [19] V. GIRAULT AND P. RAVIART, *Finite Element Methods For Navier–Stokes Equations*, Springer, Berlin, 1986.
- [20] C. GRANDMONT, *Existence and uniqueness for a two-dimensional steady-state fluid-structure interaction problem*, C. R. Acad. Sci. Paris Sér. I Math., 326 (1998), pp. 651–656.
- [21] C. GRANDMONT AND Y. MADAY, *Existence for an unsteady fluid-structure interaction problem*, M2AN Math. Model. Numer. Anal., 34 (2000), pp. 609–636.
- [22] C. GRANDMONT AND Y. MADAY, *Existence of solutions for a two-dimensional unsteady fluid-structure interaction problem*, C. R. Acad. Sci. Paris Sér. I Math., 326 (1998), pp. 525–530.
- [23] M. D. GUNZBURGER AND L. S. HOU, *Treating inhomogeneous essential boundary conditions in finite element methods and the calculation of boundary stresses*, SIAM J. Numer. Anal., 29 (1992), pp. 390–424.
- [24] P. GRISVARD, *Elliptic Problems in Nonsmooth Domains*, Pitman, Boston, 1985.
- [25] L. HOU, *Error estimates for semidiscrete finite element approximations of the Stokes equations under minimal regularity assumptions*, J. Sci. Comput., 16 (2001), pp. 287–317.
- [26] G. HSIAO, R. KLEINMAN AND G. ROACH, *Weak solutions of fluid-solid interaction problems*, Math. Nachr., 218 (2000), pp.139–163.
- [27] P. LETALLEC AND S. MANI, *Numerical analysis of a linearized fluid-structure interaction problem*, Numer. Math., 87 (2000), pp. 317–354.
- [28] H. MORAND AND R. OHAYON, *Fluid Structure Interaction: Applied Numerical Methods*, Wiley, New York, 1995.
- [29] S. MICU AND E. ZUAZUA, *Asymptotics for the spectrum of a fluid/structure hybrid system arising in the control of noise*, SIAM J. Math. Anal., 29 (1998), pp. 967–1001.
- [30] R. OHAYON AND C. FELIPPA, EDS., *Special Issue on Fluid-Structure Interactions*, Comput. Methods Appl. Mech. Engrg., 190 (2001), pp. 2977–3292.
- [31] P.D. PANAGIOTOPOULOS, *Inequality Problems in Mechanics and Applications*, Birkhäuser Boston, Cambridge, MA, 1985.
- [32] R. RODRÍGUEZ AND J. SOLOMIN, *The order of convergence of eigenfrequencies in finite element approximations of fluid-structure interaction problems*, Math. Comp., 65 (1996), pp. 1463–1475.
- [33] M. RUMPF, *On equilibria in the interaction of fluids and elastic solids. Theory of the Navier-Stokes equations*, Ser. Adv. Math. Appl. Sci., 47 (1998), pp. 136–158.
- [34] R. SCHULKES, *Interactions of an elastic solid with a viscous fluid: Eigenmode analysis*, J. Comput. Phys., 100 (1992), pp. 270–283.